

Received Date: 18-Aug-2015

Revised Date: 23-Feb-2016

Accepted Date: 04-Mar-2016

Article Type: Articles

Relatively stable response of fruiting stage to warming and cooling relative to other phenological events

Short title: Stable fruit set under warming and cooling

L.L Jiang^{1,4*}, S.P. Wang^{1,3,4}, F.D. Meng^{1,5}, J.C. Duan⁶, H.S. Niu⁵, G.P. Xu⁷, X.X. Zhu^{2,4}, Z.H. Zhang², C.Y. Luo², S.J. Cui^{1,4}, Y.M. Li¹, X.E. Li¹, Q. Wang^{1,5}, Y. Zhou^{1,5}, X.Y. Bao⁵, Y.N. Li², T. Dorji^{1,4}, S.L. Piao¹, P. Ciais⁸, J. Peñuelas^{9,10}, M.Y. Du¹¹, X.Q. Zhao², L. Zhao², F.W. Zhang², Guoji Wang¹²

1. Key Laboratory of Alpine Ecology and Biodiversity, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China.

2. Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining 810008, China.

3. CAS Center for Excellence in Tibetan Plateau Earth Science of the Chinese Academy of Sciences, Beijing 100101, China

4. Naqu Integrated Observation and Research Station of Ecology and Environment, Tibet University and Institute of Tibetan Plateau Research of the Chinese Academy of Sciences,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ecy.1450

This article is protected by copyright. All rights reserved.

Lasa 850012, China

5. University of Chinese Academy of Sciences, Beijing 100049, China.

6. Binhai Research Institute in Tianjin, Tianjin 300457, China.

7. Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, Guangxi 541006, China.

8. Laboratoire des Sciences du Climat et de l'Environnement, CEA CNRS UVSQ, 91191 Gif sur Yvette, France.

9. CREAM, Cerdanyola del Valles, Barcelona 08193, Catalonia, Spain.

10. CSIC, Global Ecology Unit CREAM-CEAB-CSIC-UAB, Cerdanyola del Vallès, Barcelona 08193, Catalonia, Spain.

11. National Institute for Agro-Environment Sciences, Tsukuba, 305-8604, Japan.

12. Oregon State University Agriculture and Natural Resource Program at Eastern Oregon University, La Grande, OR 97850, USA

Corresponding authors:

L.L. Jiang, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China. Cell phone: 86-13301321910; E-mail: lljiang@itpcas.ac.cn

L.L. Jiang, S.P. Wang, F.D. Meng, J.C. Duan and H.S. Niu were equal contribution to the paper.

This article is protected by copyright. All rights reserved.

Abstract The timing of the fruit-set stage (i.e. start and end of fruit set) is crucial in a plant's life cycle, but its response to temperature change is still unclear. We investigated the timing of seven phenological events, including fruit-set dates during 3 years for six alpine plants transplanted to warmer (+ ~3.5 °C in soils) and cooler (- ~3.5 °C in soils) locations along an altitudinal gradient in the Tibetan area. We found that fruit-set dates remained relatively stable under both warming and cooling during the 3-years transplant experiment. Three earlier phenological events (emergence of first leaf, first bud set, and first flowering) and two later phenological events (first leaf coloring and complete leaf coloring) were earlier by 4.8-8.2 days °C⁻¹ and later by 3.2-7.1 days °C⁻¹ in response to warming. Conversely, cooling delayed the three earlier events by 3.8-6.9 days °C⁻¹ and advanced the two later events by 3.2-8.1 days °C⁻¹ for all plant species. The timing of the first and/or last fruit-set dates, however, did not change significantly compared to earlier and later phenological events. Statistical analyses also showed that the dates of fruit set were not significantly correlated or had lower correlations with changes of soil temperature relative to the earlier and later phenological events. Alpine plants may thus acclimate to changes in temperature for their fruiting function by maintaining relatively stable timings of fruit set compared with other phenological events to maximize the success of seed maturation and dispersal in response to short-term warming or cooling.

Keywords: phenological sequence; seed-production stage; alpine plants; early-spring flowering plants; mid-summer flowering plants; temperature change; Tibetan plateau

Introduction

Most studies of the response of phenological events to temperature have been based on the analysis of events early or late in the season such as leaf onset, first flowering, and senescence (Walker et al. 2006, Amano et al. 2010, Yu et al. 2010; Shen et al. 2011; Piao et al. 2011; Cook et al. 2012; Wolkovich et al. 2012; Wang et al. 2014a). Plants may face trade-offs between the adjustment of one phenological event to temperature and the timing of subsequent events during the growing season (Post et al. 2008; Sherry et al. 2007, 2011; Haggerty and Galloway 2011; Dorji et al., 2013; CaraDonna et al. 2014; Wang et al. 2014b). This is illustrated by Fig. 1. Changes of flowering dates may alter the timing of fruit maturation, whereas fruit phenology in turn determines seed maturation and dispersal, which further feeds back on the diversity of species in an ecosystem (Primack 1987). The response of timing of fruiting to warming relative to other phenological events is still unclear because of inconsistent results from different plant species (Post et al. 2008; Haggerty and Galloway 2011; Sherry et al. 2007) and because few studies have analyzed the effects from both warming and cooling at the same location.

The timing of fruit set is closely related to production of offspring, survival and reproductive success (Primack 1987; Miller-Rushing and Primack 2008). The response of timing of fruit set to changes in temperature in alpine grassland communities has received little attention, either from direct observations or manipulative experiments (Dorji et al. 2013; Wang et al. 2014b). Temperate regions have a fruiting period that generally occurs in late summer or in autumn (Ting et al. 2008). For alpine communities where the growing season is short, we hypothesize that the fruiting stage does not change in response to short-term warming or

Accepted Article

cooling. Fruit set should indeed occur during a narrow period of the warmer days of the year due to large diurnal variations of temperature early and late in the growing season in mountainous regions (Li et al. 2004), which causes a risk of reproduction failure for species setting seeds too early or too late. This area of research has provided scant data. We thus tested the hypothesis that alpine plants maintain a relatively stable fruit-set date because they will be more exposed to reproduction failure if they immediately adjust the timing of their fruit production to a short-term temperature change. We tested this hypothesis in a high-altitude meadow on the Tibetan plateau. The plateau has warmed considerably over the last 40 years but also has years cooler than usual (Li et al. 2004) and is thus a suitable location for obtaining an understanding of how fruiting responds to temperature changes for projecting future plant diversity and ecosystemic productivity. Short-term warming and cooling occur on decadal and inter-annual time scales as part of natural climatic variability. Several cooling events occurred during the Holocene (Mayle and Cwynar 1995) and the last century, e.g. after volcanic eruptions (Sirocko et al. 2012), but the phenological responses to cooling remain unclear (Wang et al. 2014a) because most experiments that manipulate ecosystems have focused on warming rather than cooling. In previous study, we already showed that looking at warming alone over- or under-estimated the sensitivity of flowering date to temperature change if one does not compare the response from warming experiment to the one of independent cooling experiments (Wang et al. 2014a). We thus conducted a reciprocal transplantation experiment to expose grassland communities to both warming and cooling. The transplanted blocks were monitored from 2008 to 2010 (Wang et al. 2014b) to evaluate the response of phenological events (Fig. 1) to warming and cooling. The aims of

This article is protected by copyright. All rights reserved.

Accepted Article

the study were to determine the changes in the dates of the first fruit set (FFS) and the last fruit set (LFS) in response to warming and cooling and to compare these changes to temporal shifts in early-season (the emergence of first leaf (EFL), first budding-set (FBS), and first flowering (FF)) and late-season (the first leaf coloring (FLC) and complete leaf coloring (CLC)) phenological events.

Materials and methods

Experimental site and data collection

Details of the experimental site and design and the monitoring of the soil temperature, soil moisture, and phenological events are reported by Wang et al. (2014a, b). In brief, the experiment was conducted at the Haibei Alpine Meadow Ecosystem Research Station of the Chinese Academy of Sciences in Qinghai China (37°37'N, 101°12'E) along a 3200-3800 m altitudinal gradient (3200, 3400, 3600, and 3800 m). Four plant communities within 9 km of each other were selected for study. Twelve intact soil blocks (100×100 cm and 30-40 cm deep) (i.e. 30 cm depth at 3800 m due to the shallower soil layer) with attached vegetation from each altitude were reciprocally transplanted across the altitudinal gradient after the soils started to thaw in early May 2007 (Wang et al. 2014b). Three replicates were transplanted from each altitude, and these intact soil blocks were fully randomized throughout the study site.

Measurements of soil temperature, soil moisture, and phenological events

HOBO weather stations (Onset Computer Corporation, Cape Cod, USA) were established at the center of each site to monitor soil temperature (ST) at 5-cm depths with 1-minute

This article is protected by copyright. All rights reserved.

Accepted Article

intervals, and the data were stored in a data logger. Annual average soil temperatures at 5-cm soil depths for 2008-2010 were 3.9, 2.5, 2.0, and 0.4 °C at 3200, 3400, 3600, and 3800 m, respectively (Wang et al. 2014a).

Six common plant species from these blocks were monitored for all phenological events at each elevation during the growing seasons of 2008-2010. Events were monitored for six representative species at each elevation at intervals of 4-5 days from early April to mid-November in each year. The six species were two early-season flowering sedges *Kobresia humilis* (Kh) and *Carex scabrirostris* (Cs), two mid-season flowering forbs *Potentilla anserine* (Pa) and *P. nivea* (Pn), and two mid-season flowering grasses *Poa pratensis* (Pp) and *Stipa aliena* (Sa). Ten individuals of each forb and ten stems of each graminoid in each plot along the gradient were marked during the previous autumn so that phenological events affecting individual plants could be accurately monitored throughout the growing season. Data were collected at intervals of 4-5, The onset dates of seven phenological events were recorded (Wang et al. 2014b). No data were obtained at 3600 m in 2010 because mice destroyed the plots.

Data analysis

An analysis of variance (ANOVA) and multi-comparisons were conducted with SPSS version 22.0 (SPSS Inc., Chicago, USA), and the data were fitted with three types of functions using R version 3.1.0. Linear mixed models with repeated measurements were used for the ANOVA. Type III sum of squares was adopted for the changes in the start date compared to the donor site due to the missing data at 3600 m in 2010. For the warming and cooling due to the transplantings, a plot (i.e. a soil block) was a subject, the original (i.e. donor) and

This article is protected by copyright. All rights reserved.

transplantation (i.e. receptor) sites were between-subject factors, and species and year were within-subject factors. The changes of start date under warming and cooling were analyzed by one-way ANOVAs, and multi-comparisons of all variables were conducted using least significant difference (LSD) tests. Simple linear regression was performed between the average start date of each phenological event and the pooled data for the changes of soil temperatures at the receptor and donor sites due to warming and cooling to test the responses of the start date to the temperature change. Simple correlations were performed between the start date of each phenological event. The correlations were not corrected, and a p -value was added to each Pearson correlation coefficient. Significant differences are reported in the text at $p < 0.05$.

The response of FFD to the change in soil temperature was fitted with three types of models: linear, exponential, and piecewise linear (segmented) regression (Wang et al. 2014a; Iler et al. 2013). The fit of the models was compared based on Akaike's information criterion (AIC) (Iler et al. 2013). AIC values were used to determine whether a linear or nonlinear model was a better fit to the dates of phenological events. Only models with $\Delta AIC > 2$ were considered to be different (i.e. asymmetric, unequal responses to warming vs. cooling), whereas models with $\Delta AIC < 2$ were considered to have equal fits to the data (i.e. symmetric, equal responses to warming vs. cooling) (Wang et al. 2014a; Iler et al. 2013).

Results

The statistical power of the transplant experiment was evaluated by fitting the data to a full linear model (Appendix S1: Table S1). Various sources of error exist, but most items of the

This article is protected by copyright. All rights reserved.

Accepted Article

full model were significant at $p < 0.001$, with the exception of the year factor and the 5-way interaction (Appendix S1: Table S1). The response of phenological events to warming and cooling were not always the same in direction or magnitude (Fig. 2). The similarity or a difference in the shift of a given phenological event to temperature change was evaluated based on shift direction and magnitude. First, if shift directions were the same, then we compared shift magnitude. Finding the same shift magnitude of different phenological events cannot tease apart whether these events were dependent or independent on each other. Different shift directions lead us to assume that events were independent of each other.

Our results indicated that shifts for early-season phenological events under warming and cooling were of the same magnitude for the same species. For example, warming significantly advanced the average date of the three early-season events by the same magnitudes for the same species (average advances of 4.8-8.2 day °C⁻¹) and delayed the date of the two late-season events (average delays of 3.2-7.1 days °C⁻¹) (Fig. 2). In contrast, cooling delayed the three early-season events by an average of 3.8-6.9 day °C⁻¹ and advanced the two late-season events by an average of 3.2-7.1 days °C⁻¹ (Fig. 2).

By contrast, we found that the responses of FFS under warming and of LFS under cooling were relatively stable relative to the early- and late-season phenological events. The small change of the average FFS was even significantly smaller for the mid-season flowering plants (i.e. Pa, Pn, Pp, and Sa) under warming and for Pa and Pn under cooling relative to the early- and late-season phenological events, suggesting that the change of FFS was independent of the early- and late-season phenological events (Fig. 2). The magnitude of the delay for FFS,

This article is protected by copyright. All rights reserved.

however, remained approximately the same for the early-flowering plants (i.e. Kh and Cs) relative to the late-season phenological events under warming (Fig. 2). This suggests that FFS was only independent of the early-season events but not of the late-season events. The change of LFS was significantly smaller for all plant species under cooling relative to the early- and late-season phenological events (Fig. 2).

We observed asymmetric responses for the phenological events to warming and cooling in most situations. Most of phenological events for all species responded asymmetrically to warming and cooling, as indicated by the difference of Akaike's information criterion (ΔAIC) between the three fitted models, because ΔAIC was <2 for the three models in most situations (Table 1).

Correlations were determined between dates of phenological events and changes in soil temperature by pooling warming and cooling data together (Fig. 3). The three early-season events, namely the emergence of the first leaf (EFL), first bud set (FBS), and first flowering (FF), were negatively correlated for all species with the difference in soil temperature between the receptor and donor sites (ΔT) (Fig. 3 and Appendix S2: Table S1). Leaf senescence late-season events, namely the dates of first leaf coloring (FLC) and complete leaf coloring (CLC), were on the other hand positively correlated with ΔT (Fig. 3 and Appendix S2: Table S1). These correlations indicated that warming lengthened the growing season both by advancing onset and delaying senescence. Neither FFS nor LFS, however, were significantly correlated with ΔT for either the early-flowering (Kh and Cs) or the mid-season flowering forbs (Pa and Pn) (Fig. 3A-3D and Appendix S2: Table S1). FFS and LFS were significantly correlated with ΔT for the two other mid-season flowering grasses (i.e. Pp and Pq).

This article is protected by copyright. All rights reserved.

Sa) (Fig. 3 and Appendix S2: Table S1), but the slopes of the regression lines were 2-5 fold smaller than those obtained for the early and late-season events (Appendix S2: Table S1). In sum, the timing of fruiting events remained relatively stable during three years of warming and cooling.

Discussion

The reciprocal transplants may have experienced changes in soil moisture along the elevational gradient, but previous studies have indicated that alpine-plant phenology is primarily affected by changes in temperature, not moisture (Wang et al. 2014a, b). We monitored only six species, but our results suggested that their fruit-set events in the community did not change in response to warming and cooling but that the other phenological events late and early in the season were relatively responsive to temperature change. Fruiting was found to advance more than budding and flowering events during warming experiments performed in temperate grasslands (Post et al., 2008; Haggerty and Galloway, 2011). Sherry et al. (2007) reported that changes in fruiting under warming were species-specific, depending on their specific flowering dates. These results, compared with ours for the alpine meadow of Tibet led us to conclude that fruiting may not respond uniformly to climate change across different grassland ecosystems, given plant phylogenetics and the biotic interactions within communities (Cortés-Flores et al. 2013). Fruiting time may depend on trade-offs between resource availability, the presence of pollinators, the abundance of herbivores, suitable conditions for seed production, and phylogenetic constraints (Murali and Sukumar 1994; Wright and Calderón 1995; Sherry et al. 2007; Nakajima et al. 2012).

This article is protected by copyright. All rights reserved.

The relatively stable fruiting date in response to short-term changes in soil temperature for the Tibetan meadows in this study can be attributed to three potential mechanisms. Firstly, similar to first flowering events studied in a previous report (Wang et al. 2014a), the response of fruiting to warming and cooling differed in direction and magnitude (Fig. 2), implying that short term cooling (e.g. occurrence of cold years) could modify the effective magnitude of long-term warming on fruiting because of the asymmetric annual response to warming and cooling (Table 1). For example, for the early-flowering plants (i.e. Kh and Cs), the change of FFS was correlated negatively with ΔT under cooling but positively with ΔT under warming (Fig. S1), which could explain the lack of significant correlation between ΔFFS and ΔT when pooled warming and cooling data together (Fig. 3 and Appendix S2: Table S1). The asymmetric response of LFS to warming and cooling for the early-flowering plants may have a similar explanation. In contrast, the changes of FFS and LFS were not correlated with ΔT under warming for the mid-season flowering grasses (i.e. Pp and Sa), but FFS and LFS were correlated with ΔT when separating the warming and cooling data (Fig. 3, Appendix S2: Table S1, Appendix S4: Fig. S1, and Appendix S5: Fig. S1). Similar to first flowering (Wang et al. 2014a), warming experiments alone may thus over- or underestimate the response of fruit set, depending on species, relative to warming and cooling experiments. The effect of cooler years which can occur as part of the natural climate variability at each site should thus be measured when evaluating the effects of warming on phenology, because the alternation of cool and warm years during a decade with a warming trend will not cause the same response as a continuous warming trend.

Secondly, each life-history stage of a plant is to some extent dependent on or constrained by the preceding and succeeding stages (Sherry et al. 2007; Post et al. 2008; Wang et al. 2014b). Plant phylogenetic constraints in the fruit-set stage severely limit adaptive potential (Bradshaw 1984). Previous studies have shown that the temporal patterns of fruiting can depend on flowering phenology (Primack 1987). We, however, found that the changes of flowering and fruiting dates were poorly correlated for warming ($r^2=0.16$) and warming+cooling ($r^2=0.19$), even though their relationships were significant (Appendix S3, Table S1), and well correlated only for cooling alone ($r^2=0.60$) (Appendix S3, Table S1). The change of FF was not correlated with the change of LFS under warming, and the correlations were low under cooling ($r^2=0.21$) and under warming+cooling ($r^2=0.04$) (Appendix S3, Table S1). These results suggest that the adaptive adjustment of fruiting time may be independent on flowering phenology (Thompson and Willson 1979; Stiles 1980, Eriksson and Ehrlén 1991). Moreover, fruiting patterns were the same for the two plant species of each family (i.e. Kh and Cs of the Cyperaceae family, Pa and Pn of the Asteraceae family, and Pp and Sa of the Gramineae family). Flowering usually simply occurs when a plant reaches a critical size or developmental stage or accumulated degree days (Diekmann 1996; Wang et al. 2014a), whereas the phenology of fruit set is governed by different constraints (Jordano 1992). Fruiting time, by definition, equals flowering time plus the developmental time from flower to fruit. One likely constraint on this developmental time is seed size, which is significantly correlated with developmental time (Eriksson and Ehrlén 1991), although fruit production is largely controlled by the accumulation of photosynthates (French 1992). A relatively stable or delayed timing of fruit set could thus be associated with the formation of larger seeds

This article is protected by copyright. All rights reserved.

(Primack 1987; Sherry et al. 2007; Menzel et al. 2011), which could improve seedling establishment and growth (Wang et al. 2012), and not limit the period of seed maturation, especially for the early-flowering plants in the warming environments. Late-flowering individuals, however, may not have enough time for fruit maturation (Helgen and Barrett 1987; Rathcke 1988; Kudo 1993; Wang et al. 2014b), which would be an important constraint. Minor shifts to the end of fruit set under cooling (Fig. 2) could thus mainly provide plants under cooling in alpine regions with enough time for seed maturation due to a shorter vegetative stage after fruit set (Kudo 1997).

Thirdly, biotic interactions could also be important drivers in determining the fruiting stage. Herbivores exert a strong selection pressure on plants in grasslands. For example, earlier-flowering plants (e.g. Kh and Cs) (Fig. 2) are grazed by animals such as sheep, yaks, mice, and caterpillars (Zhao 2009; Cao et al. 2015). A possible strategy of plants in a community for reducing herbivorous pressure is a short period of synchronous production of leaves, buds, and flowers, thereby satiating the herbivores (Aide 1991; Fenner 1998).

Early-flowering plants complete their floral primordia the previous autumn and winter (Körner 2003), and limited accumulation of photosynthates can support their reproductive development. Their later FFS under warming and cooling (Fig. 2), however, may reduce the loss of seeds, because breeding birds need the number of highly nutritious seeds to peak during the early spring. The seeds would benefit from 'temporal dispersal' with a persistent seed bank because of the lower expected flight distances of birds during the breeding season.

The differentiation of floral primordia in mid-season-flowering plants is synchronized with vegetative growth before flowering (Körner 2003), and vegetative growth before budding is

This article is protected by copyright. All rights reserved.

slow (Zhou 2001). In particular, mid-season-flowering forbs (i.e. Pa and Pn) with large yellow flowers are under the vegetative canopy of insect-pollinated plants. Simultaneous phenological sequences (Fig. 2), especially joint floral display, may thus increase visitation rates, leading to enhanced seed set for one or both species (Thompson 1978, 1981, 1982; Schemske 1981), because synchronous blooming would attract proportionally more pollinators than asynchronous blooming (Rathcke 1983). Moreover, earlier flowering but a stable fruiting date (i.e. longer period of flowering) under warming should increase pollinator activity in a situation when plants and pollinators respond to warming differently (Kudo and Suzuki 2002) to avoid pollination deficits due to phenological mismatches between plants and pollinators (Kudo et al. 2004; Memmott et al. 2007; Williams and Jackson 2007). Neither warming nor cooling altered or reduced shifts in the fruit-set stage in the taller mid-season-flowering grasses in the community (Pp and Sa), probably because most bird-dispersed plants ripen their fruit during the peak of bird abundance, usually in early fall (Stiles 1980), to escape pathogens and non-disperser frugivores (Thompson and Willson 1979). Many temperate regions receive massive annual influxes of migratory frugivorous birds (Herrera 1998; Teller'a and Pérez-Tris 2003, 2007). Their abundance thus varies seasonally, peaking in autumn during migration to winter quarters, which may facilitate the spread of seeds over larger distances (Snow 1971; Thompson and Willson 1979; Herrera 1984).

Conclusions

Our finding of a relatively stable fruiting stage is analogous to the earlier arrival of migratory birds at their breeding grounds with warming (Root et al. 2003) but with a comparatively minor shift in egg-laying (Both and Visser 2005). Changes in the commencement of flowering are likely to lead to changes in fruit maturation and seed dispersal under warming and cooling periods in alpine regions. Fruit production is the central function of flowering, and seed dissemination is the major function of the fruit, so we should expect relatively stable fruiting phenology to be influenced by selective pressures that would favor successful seed maturation and dispersal. Information for fruiting phenology is of central importance for the conservation of community diversity and is especially valuable for alpine regions that support a diversity of fruit-dependent species.

Acknowledgements

This work was supported by projects from the National Science Foundation of China (41230750), National Basic Research Program (2013CB956000), Strategic Priority Research Program (B) of the Chinese Academy of Sciences (XDB03030403), and National Science Foundation of China (31272488 and 31470524) and China postdoctoral science foundation (2013M541050). Philippe Ciais and Josep Peñuelas acknowledge support from the European Research Council through Synergy grant ERC-2013-SyG-610028 “IMBALANCE-P”

References

- Aide, T. M. 1991. Synchronous leaf production and herbivory in juveniles of *Gustavia superba*. *Oecologia* 88:511-514.
- Amano, T., R. J. Smithers, T. H. Sparks and W. J. A. Sutherland. 2010. 250-year index of first flowering dates and its response to temperature changes. *Proceeding of Royal Society London B* 277:2451-2457.
- Both, C. and M. E. Visser. 2005. The effect of climate change on the correlation between avian life-history traits. *Global Change Biology* 11:1606-1613.
- Bradshaw, A. D. 1984. The importance of evolutionary ideas in ecology – and vice versa. *Evolutionary Ecology*, ed B. Shorrocks (Symposium BES 23, Blackwells Scientific Publications, Oxford), pp 1-25.
- Cao, H., X. Q. Zhao, S. P. Wang, L. Zhao, J. C. Duan, Z. H. Zhang, S. D. Ge and X. X. Zhu. 2015. Grazing intensifies degradation of a Tibetan Plateau alpine meadow through plant-pest interaction. *Ecology and Evolution* doi:10.1002/ece3.1537.
- CaraDonna, P. J., A. M. Iler and D. W. Inouye. 2014. Shifts in flowering phenology reshape a subalpine plant community. *Proceedings of the National Academy of Sciences* 111:4916-4921.
- Cook, B. I., E. M. Wolkovich and C. Parmesan. 2012. Divergent responses to spring and winter warming drive community level flowering trends. *Proceedings of the National Academy of Sciences* 109:9000-9005.
- Cortés-Flores, J., E. Andresen, G. Cornejo-Tenorio and G. Ibarra-Manríquez. 2013. Fruiting phenology of seed dispersal syndromes in a Mexican Neotropical temperate forest. *Forest*
- This article is protected by copyright. All rights reserved.

Ecological Manage, 289, 445-454.

- Diekmann, M. 1996. Relationship between flowering phenology of perennial herbs and meteorological data in deciduous forests of Sweden. *Canadian Journal of Botany* 74:528-537.
- Dorji, T., Ø. Totland, S. Moe, K. A. Hopping, J. B. Pan and J. A. Klein. 2013. Plant functional traits mediate reproductive phenology and success in response to experimental warming and snow addition in Tibet. *Global Change Biology* 19:459-472.
- Eriksson, O. and J. Ehrlén. 1991. Phenological variation in fruit characteristics in vertebrate-dispersed plants. *Oecologia* 86:463-470.
- Fenner, M. 1998. The phenology of growth and reproduction in plants. *Perspective of Plant Ecology and Evolution System* 1:78-91.
- French, K. 1992. Phenology of fleshy fruits in a wet sclerophyll forest in southeastern Australia: Are birds an important influence? *Oecologia* 90:366-373.
- Haggerty, B. P. and L. F. Galloway. 2011. Response of individual components of reproductive phenology to growing season length in a monocarpic herb. *Journal of Ecology* 99:242-253.
- Helenurm, K. and S. C.H. Barrett. 1987. The reproductive biology of boreal forest herbs. 2. Phenology of flowering and fruiting. *Canadian Journal of Botany* 65:2047-2056.
- Herrera, C. M. 1984. A study of avian frugivores, bird-dispersed plants, and their interactions in Mediterranean shrubland. *Ecological Monograph* 54:1-23.
- Herrera, C. M. 1998. Long-term dynamics of Mediterranean frugivorous birds and fleshy-fruits: a 12-year study. *Ecological Monograph* 68:511-538.

- Iler, A. M., T. T. Hoye, D. W. Inouye and N. M. Schmidt. 2013 Nonlinear flowering responses to climate: are species approaching their limits of phenological change? *Philosophical Transactions of Royal Society B* 368:10.1098/rstb.2012.0489
- Jordano, P. 1992. Fruits and frugivory. *Seeds: The Ecology of Regeneration in Plant Communities*, ed. M. Fenner (CAB International, Oxford), pp 105-156.
- Körner, C. (ed). 2003. *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems*. Springer-Verlag, New York (ed. 2). pp 260-262.
- Kudo, G. 1993. Relationship between flowering time and fruit set of the entomophilous alpine shrub, *Rhododendron aureum* (Ericaceae), inhabiting snow patches. *American Journal of Botany* 80:1300-1304.
- Kudo, G. 1997. Sex expression and fruit set of an andromonoecious herb, *Peucedanum multivittatum* (Umbelliferae) along a snowmelt gradient. *Opera Botany* 132:121-128.
- Kudo, G. and S. Suzuki. 2002. Relationships between flowering phenology and fruit-set of dwarf shrubs in alpine fellfield in Northern Japan: a comparison with a subarctic heathland in northern Sweden. *Arctic, Antarctic and Alpine Research* 34:185-190.
- Kudo, G., Y. Nishikawa, I. Kasagi and S. Kosuge. 2004. Does seed production of spring ephemerals decrease when spring comes early? *Ecological Research* 19:255-259.
- Li, Y., X. Zhao, G. Cao, L. Zhao and Q. Wang. 2004. Analyses on climates and vegetation productivity background at Haibei alpine meadow ecosystem research station. *Plateau Meteorology* 23:558-567.
- Mayle, F. E. and L. C. Cwynar. 1995. Impact of the Younger Dryas cooling event upon lowland vegetation of Maritime Canada. *Ecological Monograph* 65:129-154.
- This article is protected by copyright. All rights reserved.

- Menzel, A., H. Seifert and N. Estrella. 2011. Effects of recent warm and cold spells on European plant phenology. *International Journal of Biometeorology* 55:921-932.
- Memmott, J., P. G. Craze, N. M. Waser and M. V. Price. 2007. Global warming and the disruption of plant–pollinator interactions. *Ecology Letters* 10:710-717.
- Miller-Rushing, A. J. and R. B. Primack. 2008. Global warming and flowering times in Thoreau’s Concord: a community perspective. *Ecology* 89:332-341.
- Murali, K. S. and R. Sukumar. 1994. Reproductive phenology of a tropical dry forest in Mudumalai, southern India. *Journal of Ecology* 82:759-767.
- Nakajima, A. et al. 2012. Spatial and elevational variation in fruiting phenology of a deciduous oak (*Quercus crispula*) and its effect on foraging behavior of the Asiatic black bear (*Ursus thibetanus*). *Ecological Research* 27:529-538.
- Piao, S., M. Cui, A. Chen, X. Wang, P. Ciais, J. Liu and Y. Tang. 2011. Altitude and temperature dependence of change in the spring vegetation green-up date from 1982 to 2006 in the Qinghai-Xizang Plateau. *Agriculture, Forest Meteorology* 151:1599-1608.
- Post, E. S., C. Pedersen, C. C. Wilmer and M. C. Forchhammer. 2008. Phenological sequences reveal aggregate life history response to climatic warming. *Ecology* 89:363-370.
- Primack, R. B. 1987. Relationships among flowers, fruits, and seeds. *Annual Review of Ecology and System* 18:409-430.
- Rathcke, B. J. 1983. Competition and facilitation among plants for pollination. *Pollination Biology*, ed. L.A. Real (Academic Press, New York), pp 305-338.
- Rathcke, B. J. 1988. Flowering phenologies in a shrub community: competition and
- This article is protected by copyright. All rights reserved.

constraints. *Journal of Ecology* 76:975-994.

Root, T. L., J. T. Price, K. R. Hall, S. H. Schneider, C. Rosenzweig and J. A. Pounds. 2003.

Fingerprints of global warming on wild animals and plants. *Nature* 421:57-60.

Schemske, D. W. 1981. Floral convergence and pollinator sharing in two bee-pollinated tropical herbs. *Ecology* 62:946-954.

Shen, M., Y. Tang, J. Chen, X. Zhu and Y. Zheng. 2011. Influences of temperature and precipitation before the growing season on spring phenology in grasslands of the central and eastern Qinghai-Tibetan Plateau. *Agriculture, Forest Meteorology* 151:1711-1722.

Sherry, R. A. et al. 2007. Divergence of reproductive phenology under climate warming.

Proceedings of the National Academy of Sciences 104:198-202.

Sherry, R. A. et al. 2011. Changes in duration of reproductive phases and lagged phenological response to experimental climate warming. *Plant Ecology and Diversity* 4:23-35.

Sirocko, F., H. Brunck and S. Pfahl. 2012. Solar influence on winter severity in central Europe. *Geography Research Letters* 39:L16704, doi:10.1029/2012GL052412.

Snow, D. W. 1971. Evolutionary aspects of fruit-eating by birds. *Ibis* 113:194-202.

Stiles, E. W. 1980. Patterns of fruit presentation and seed dispersal in bird-disseminated woody plants in the eastern deciduous forest. *American Naturalist* 116:670-688.

Tellería, J. L. and J. Pérez-Tris. 2003. Seasonal distribution of a migratory bird: effects of local and regional resource tracking. *Journal of Biogeography* 30:1583-1591.

Tellería, J. L. and J. Pérez-Tris. 2007. Habitat effects on resource tracking ability: do wintering blackcaps *Sylvia atricapilla* track fruit availability? *Ibis* 149:18-25.

Thompson, J. D. 1978. Effect of stand composition on insect visitation in two-species

This article is protected by copyright. All rights reserved.

Accepted Article

mixtures of Hieracium. *American Midland Naturelist* 100:431-440.

Thompson, J. N. and M. F. Willson. 1979. Evolution of temperate fruit/bird interactions: phenological strategies. *Evolution* 33:973-982.

Thompson, J. D. 1981. Spatial and temporal components of resource assessment of flower-feeding insects. *Journal of Animal Ecology* 50:49-59.

Thompson, J. D. 1982. Pattern of visitation by animal pollinators. *Oikos*, 39, 241-250.

Ting S, Hartley S, Burns KC (2008) Global patterns in fruiting seasons. *Global Ecology and Biogeography* 17:648-657.

Walker, M. D. et al. 2006. Plant community responses to experimental warming across the tundra biome. *Proceedings of the National Academy of Sciences* 103:1342-1346.

Wang, Q. J. and S. B. Shi. 2001. Biomass and its forming mechanism. *Kobresia meadow in China*, ed Zhou XM (Beijing: Science Press), pp 134.

Wang, S. P. et al. 2012. Effects of warming and grazing on soil N availability, species composition and ANPP in alpine meadow. *Ecology* 93:2365-2376.

Wang, S. P. et al. 2014a. Asymmetric sensitivity of first flowering date to warming and cooling in alpine plants. *Ecology* 95:3387-3398.

Wang, S. P. et al. 2014b. Timing and duration of phenological sequences of alpine plants along an elevation gradient on the Tibetan plateau. *Agriculture, Forest Meteorology* 189-190:220-228.

Warton, D. I. et al. 2012. SMATR 3 - an R package for estimation and inference about allometric lines. *Methods in Ecology and Evolution* 3:257-259.

Williams, J. W. and S. T. Jackson. 2007. Novel climates, no-analog communities, and

This article is protected by copyright. All rights reserved.

ecological surprises. *Frontier of Ecology and Environment* 5:475-482.

Wolkovich, E. M. et al. 2012. Warming experiments underpredict plant phenological responses to climate change. *Nature* 485:494-497.

Wright, S. J. and O. Calderón. 1995. Phylogenetic patterns among tropical flowering phenologies. *Journal of Ecology* 83:937-948.

Yu, H., E. Luedeling and J. Xu. 2010. Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proceedings of the National Academy of Sciences* 107:22151-22156.

Zhao, X. Q. (ed.). 2009. *Alpine Meadow Ecosystem and Global Change*. Beijing: Science Press. pp 188.

Table 1 Akaike's information criterion of the three fitted models for various phenological events.

Phenological event	Model	Kh	Cs	Pa	Pn	Pp	Sa
EFL	Linear	526.9	518.6	492.9	541.1	509.4	507.5
	Seg.	521.4	513.7	473.2	527.4	507.8	506.9
	Exp.	527.8	519.3	482.7	535.1	508.6	506.9
FBS	Linear	594.7	588.1	659.6	583.7	620.0	568.0
	Seg.	582.7	565.1	659.6	576.3	622.9	569.5
	Exp.	584.4	566.6	659.9	579.2	621.4	569.0

FFS	Linear	609.8	645.5	607.1	606.5	595.2	505.6
	Seg.	543.5	609.8	NA	605.4	594.7	502.0
	Exp.	614.3	639.1	609.0	608.4	594.3	501.9
LFS	Linear	619.2	667.5	619.4	647.9	581.4	718.5
	Seg.	572.1	634.6	608.8	639.1	580.9	714.9
	Exp.	606.2	653.1	620.5	NA	581.9	717.7
FLC	Linear	570.5	676.9	612.9	629.7	684.4	681.5
	Seg.	NA	676.1	581.3	601.5	682.5	680.2
	Exp.	572.3	678.3	587.6	608.3	686.4	683.3
CLC	Linear	514.1	530.7	614.6	644.8	595.1	586.5
	Seg.	512.1	517.1	592.8	633.6	591.9	583.9
	Exp.	515.0	523.1	603.9	639.3	597.0	588.5

Notes: Kh, *Kobresia humilis*; Cs, *Carex scabrirostris*; Pa, *Potentilla anserine*; Pn, *P. nivea*; Pp, *Poa pratensis*; Sa, *Stipa aliena*; EFL, emergence of first leaf; FBS, first budding/booting set; FFS, first fruit set for forbs or seed set for graminoids; VAFS, vegetative stage after fruit/seed set; FLC, first leaf coloring; CLC, complete leaf coloring; Linear, linear regression; Seg., segmented regression; Exp., exponential regression; and NA, not available. Values are considered to be the same if the difference in Akaike's information criterion (AIC) is <2. The results of first flowering were reported by Wang et al. (2014a). The lowest AIC values are bolded.

Accepted Article

Fig. 1 Conceptual diagram illustrating the changes of phenological sequence with changes of soil temperature during the growing season. The black line represents no temperature change (i.e. donor site), The red line represents warming (i.e. downward translocation), and the blue line represents cooling (i.e. upward translocation). EFL, emergence of first leaf; FBS, first bud/boot set; FF, first flowering; FFS, first fruit set for forbs or seed set for graminoids; LFS, last fruit/seed set; FLC, first leaf coloring; and CLC, complete leaf coloring. We hypothesize that the seed-production (shaded area) is a relative stable under a short-term warming and cooling.

Fig. 2 Comparison of changes of start or end date of phenological events under warming or cooling. EFL, emergence of first leaf; FBS, first bud/boot set; FF, first flowering; FFS, first fruit set for forbs or seed set for graminoids; LFS, last fruit/seed set; FLC, first leaf coloring; and CLC, complete leaf coloring. Early-season phenological events include EFL, FBS, and FF. Late-season phenological events include FLC and CLC. Early-season phenological events are above the solid line. Late-season phenological events are below the dashed line. Fruit events are between the solid and dashed lines. Δ days is the difference between transplantation site (i.e. receptor site) and original site (i.e. donor site). Positive and negative values are delays and advances, respectively, of phenological events when transplanted compared to the donor site. Kh, *Kobresia humilis*; Cs, *Carex scabrirostris*; Pa, *Potentilla anserine*; Pn, *P. nivea*; Pp, *Poa pratensis*; and Sa, *Stipa aliena*. The data are means \pm SE. Different letters indicate significant differences at the 0.05 level.

Accepted Article

Fig. 3 Linear regressions between the first day of different phenological events and differences in soil temperature at a depth of 5 cm from receptor and donor sites. A, B, C, D, E, and F are *Kobresia humilis* (Kh), *Carex scabrirostris* (Cs), *Potentilla anserine* (Pa), *P. nivea* (Pn), *Poa pratensis* (Pp), and *Stipa aliena* (Sa), respectively. EFL, emergence of first leaf; FBS, first bud/boot set; FF, first flowering; FFS, first fruit set for forbs or seed-set for graminoids; LFS, last fruit/seed set; FLC, first leaf coloring; and CLC, complete leaf coloring. Early-season phenological events include EFL, FBS, and FF. Late-season phenological events include FLC and CLC. Differences in soil temperature on the x-axis are for a depth of 5 cm between receptor and donor sites (°C).





