



Regional detection of canopy nitrogen in Mediterranean forests using the spaceborne MERIS Terrestrial Chlorophyll Index

Yasmina Loozen¹, Karin T. Rebel¹, Derek Karssenber², Martin J. Wassen¹, Jordi Sardans^{3,4} and Josep Peñuelas^{3,4}, Steven M. De Jong²

5 ¹Copernicus Institute of sustainable development, Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands

²Physical geography, Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands

³CSIC, Global Ecology Unit CREAM-CSIC-UAB, 08913 Bellaterra, Catalonia, Spain

⁴CREAF, 08913 Cerdanyola del Vallès, Catalonia, Spain

Correspondence to: Yasmina Loozen (y.m.a.loozen@uu.nl)

10 **Abstract.** Canopy nitrogen (N) concentration and content are linked to several vegetation processes and canopy N concentration is a state variable in global vegetation models with coupled carbon (C) and N cycles. While there is ample C data available to constrain the models, widespread N data are lacking. Remote sensing and vegetation indices have been used to detect canopy N concentration and canopy N content at the local scale in grasslands and forests. In this paper we conducted a regional case-study analysis investigating the relationship between the Medium Resolution Imaging Spectrometer (MERIS) Terrestrial Chlorophyll Index (MTCI) time series from ESA ENVISAT at 1 km spatial resolution and both canopy N concentration (%N) and canopy N content (g m^{-2}) from a Mediterranean forests inventory in the region of Catalonia, NE of Spain. The relationships between the datasets were studied after resampling both datasets to lower spatial resolutions (20 km, 15 km, 10 km and 5 km) and at the initial higher spatial resolution of 1 km. The results at the higher spatial resolution yielded significant relationships between MTCI and both canopy N concentration and content, $r^2 = 0.32$ and $r^2 = 0.17$, respectively. We also investigated these relationships per plant functional type. While the relationship between MTCI and canopy N concentration was strongest for deciduous broadleaf and mixed plots ($r^2 = 0.25$ and $r^2 = 0.47$, respectively), the relationship between MTCI and canopy N content was strongest for evergreen needleleaf trees ($r^2 = 0.20$). At the species level, canopy N concentration was strongly related to MTCI for European Beech plots ($r^2 = 0.71$). These results present a new perspective on the application of MTCI time series for canopy N detection, ultimately leading towards the generation of canopy N maps that can be used to constrain global vegetation models.

25 **Keywords:** vegetation index, MERIS, foliar nitrogen concentration, foliar nitrogen content, plant functional types, Mediterranean forest, remote sensing

1 Introduction

30 Canopy nitrogen (N) concentration is an essential state variable in regional (Ollinger and Smith, 2005) and global vegetation models including both the carbon (C) and the N cycles (Zaehle and Friend, 2010; Smith et al., 2014). This variable has been



linked to several vegetation traits and processes, at the leaf and canopy levels. At the leaf level, leaf N concentration, which represents the leaf N status expressed as a percentage of leaf dry mass (%N, $\text{N g } 100\text{g}^{-1} \text{ DM}$), has been related to photosynthetic capacity (Evans, 1989; Reich et al., 1995; Reich et al., 1997; Reich et al., 1999; Wright et al., 2004), specific leaf area, leaf life span (Reich et al., 1999; Wright et al., 2004) and light use efficiency (Kergoat et al., 2008). Leaf N concentration expressed on
35 a leaf area basis, also called leaf N content (N g m^{-2}) has also been linked with chlorophyll content, Rubisco content (Evans, 1989) and photosynthetic capacity (Evans, 1989; Reich et al., 1995). At stand scale, canopy nitrogen concentration, which represents the leaf N concentration averaged over the stand canopy, has also been found to correlate with above ground Net Primary Productivity (NPP) (Reich, 2012), while canopy N content has been linked with the canopy light use efficiency (Green et al., 2003).

40 Given their links to many vegetation processes, leaf and canopy N variables could be used to constrain N cycle modules in global vegetation models. At the global scale, ample data is available to constrain models for the C cycle; however, data to constrain the N cycle are limited. Currently, canopy N data is not widely available and canopy N sampling campaigns are time-consuming and thus expensive tasks. Moreover, upscaling from local sampling campaign measurements represents an
45 additional limitation. In this perspective, local, regional or even global remotely sensed canopy N estimates will be a valuable addition, enabling us to collect information in a less time intensive and expensive manner than traditional on-field sampling campaigns. Such near global canopy N estimates will be beneficial as input in global vegetation models or to calibrate and validate these models.

Currently, different remote sensing techniques have been applied to detect canopy N in terrestrial vegetation. Imaging spectrometry from either airborne or spaceborne sensors coupled with different analysis methods, including partial least
50 squares regression (PLS), continuum removal, spectral unmixing or vegetation indices, has proven efficient in improving N sensing capabilities at the local scale (Smith et al., 2003; Ollinger et al., 2008; Huber et al., 2008; Martin et al., 2008; Schlerf et al., 2010; Wang et al., 2016).

Among other techniques, ratios or normalized differences of reflectance bands in the Red and Near Infrared (NIR) regions of the spectrum, the so called vegetation indices (VI) (Glenn et al., 2008), are one of the most straightforward methods for canopy
55 N detection. Combined with in situ hyperspectral devices, vegetation indices have been extensively used for leaf or canopy N detection in agricultural systems (Peñuelas et al., 1994; Filella et al., 1995; Hansen and Schjoerring, 2003; Tian et al., 2011; Schlemmer et al., 2013; Li et al., 2014). Vegetation indices have also been applied to airborne or spaceborne acquired imagery in natural environments (Ramoelo et al., 2012; Wang et al., 2016).

A particular vegetation index, the MERIS Terrestrial Chlorophyll Index (MTCI) has been proposed for detecting canopy N
60 (Clevers and Gitelson, 2013). MTCI was originally computed from three reflectance bands from the Medium Resolution Imaging Spectrometer (MERIS) aboard the European Space Agency (ESA) ENVISAT satellite at a spatial resolution of 1 km. However, it can also be obtained from other sensors' reflectance data and a similar product will be available from the ESA Sentinel-2 satellite mission (Drusch et al., 2012). It was first developed to estimate chlorophyll content (Dash and Curran, 2004, 2007). Since then, other applications of this index have been described, among which the possibility to estimate Gross



65 Primary Productivity (GPP) from natural (Harris and Dash, 2010, 2011; Boyd et al., 2012) and cultivated lands (Peng and
Gitelson, 2011). Furthermore, MTCI has been used to discriminate between C3 and C4 grasses (Foody and Dash, 2007) and
to monitor vegetation phenology at the sub-regional (Boyd et al., 2011) and continental scales (Rodriguez-Galiano et al.,
2015; Crabbe et al., 2016). Regarding canopy N detection, most applications were aimed at agricultural crops using MTCI
70 values computed from in situ hyperspectral reflectance data (Tian et al., 2011; Clevers and Gitelson, 2013; Li et al., 2014). A
few were directed towards sensing N concentration in natural environments using airborne data, e.g. in temperate forests (Wang
et al., 2016), or spaceborne data, for example in grasslands (Ramoelo et al., 2012; Ullah et al., 2012) or sub-tropical forests
(Cho et al., 2013).

Remote detection of foliage N status has been extensively studied at the leaf scale (Hansen and Schjoerring, 2003; Ferwerda et
al., 2005; Pacheco-Labrador et al., 2014; Li et al., 2014) and a few studies have investigated the processes underlying the
75 relationships between vegetation indices and canopy N. Detection of foliage N status with vegetation indices is attributed to
the strong link between foliar nitrogen and chlorophyll content (Schlemmer et al., 2013) and is often based on the NIR and
red-edge region of the spectrum, hence similar to the ones used for chlorophyll detection (Filella and Penuelas, 1994; Dash and
Curran, 2004; Clevers and Gitelson, 2013). It is not clear however how this translates at the canopy level as the mechanisms
possibly modifying the remote detection of foliage N status at the canopy scale are still not clearly understood. High correlation
80 between canopy N and both NIR reflectance and albedo has been reported in boreal forests (Ollinger et al., 2008). However,
the mechanism behind these findings is still controversial. Knyazikhin et al. (2013) argued that the observed correlation solely
resulted from canopy structural differences between broad and needleleaf forests and was thus spurious. Other authors,
although agreeing that canopy structure was a confounding factor to account for, stated that the NIR – canopy N relationship
was not necessarily spurious as plant traits have been known to covary along the leaf economic spectrum (Wright et al.,
85 2004; Ollinger et al., 2013; Townsend et al., 2013).

In this context, there are several knowledge gaps that we would like to address in this paper. First, although 1 km spatial
resolution spaceborne MTCI time series are available from the ESA, MTCI has mainly been employed to detect canopy N in
agricultural applications with in situ devices and rarely in a broader range of natural ecosystems and scales using spaceborne
data. Due to its almost global coverage, MTCI time series could be applied to estimate canopy N at a larger scale. Moreover,
90 to our knowledge limited research has been conducted to sense canopy N in Mediterranean ecosystems (Serrano et al., 2002)
and even more so in Mediterranean forests. In addition, although in a temperate forest the reflectance spectrum of individual
plant functional types (PFT) has been shown to be different (Wang et al., 2016), the relationship between MTCI and canopy
N has seldom been studied and compared between PFTs. Moreover, investigating the influence of PFTs on this relationship
might give further insight into the influence of structural effects in canopy N detection. Finally, the difference between sensing
95 canopy N concentration ($N_{[\%]}$, %N) and canopy N content ($N_{[\text{area}]}$, g m^{-2}) has rarely been investigated. The relationship between
MTCI and both of these variables has been studied separately (Clevers and Gitelson, 2013; Wang et al., 2016), but very few
analyses (Mirik et al., 2005; Ullah et al., 2012) have compared the ability to detect canopy N concentration and canopy N
content simultaneously, especially in forest ecosystems.



The objective of our study is thus to investigate the relationship between the spaceborne MTCI remote sensing product and canopy N in Mediterranean forests at the regional scale. More specifically, the relationships between MTCI and both canopy N concentration and canopy N content are investigated and compared. We then also examine these relationships per PFT and at the species level.

Detection of canopy N is often limited to local scale studies due to the spatial restrictions associated with N data acquisition in the field and treatment of high spatial resolution remote sensing imagery with limited spatial coverage (Lepine et al., 2016).

Our case-study exploits the broadly and readily available MTCI time series at 1 km spatial resolution from the ESA ENVISAT mission and combines it with canopy N data, both concentration and content, from 1075 forest plots measured between 1988 and 2001 by the Catalanian National Forest Inventory (Gracia et al., 2004). First, we develop a methodology to overcome the time discrepancy between our two sets of data. Next, both data sets are resampled to the same, lower, spatial resolutions in order to overcome the initial scale discrepancy between MTCI spatial resolution (1 km) and the size of the forest plots (6 m).

Subsequently, we analyse the relationship between MTCI and both canopy N concentration and canopy N content variables, both at the resampled and initial spatial resolutions. These relationships are then stratified according to the plots PFT. The results are presented and discussed. Finally, we address the implications for future research and draw a conclusion.

2 Material and methods

2.1 Study area

Our study area corresponds to the region of Catalonia (Fig. 1) which is located in north eastern Spain and has a spatial extent of 32,114 km² (Sardans et al., 2011). While the region is characterised by a Mediterranean climate, the presence of the Pyrenees to the northwest and the Mediterranean Sea to the east creates contrasting climate conditions with an altitudinal gradient from north to south and a continental gradient from west to east. Following this pattern, the mean annual temperature varies from 1 °C in the north to 17 °C in the south (Sardans et al., 2011). While mean annual precipitation (MAP) is 1400 mm in the Pyrenees, in the south, the MAP is lower than 350 mm (Sardans et al., 2011), leading to seasonal drought (Lana and Burgueño, 1998) and fires (González and Pukkala, 2007), impacting the vegetation (Liu et al., 2015).

2.2 Data collection

2.2.1 Canopy N data

The canopy N data used in this research was collected by the Ecological and Forestry Applications Research Centre (CREAF), Universitat Autònoma de Barcelona. The data included 2300 closed canopy forest plots sampled between 1988 and 2001 by the Catalanian National Forest Inventory (Gracia et al., 2004).

The forest plots (Fig. 1) had a minimum diameter of 6 m, which varied depending on the tree density in order to include between 15 and 25 trees with a diameter at breast height of at least 5 cm. The plots were investigated for canopy N concentration



($N_{[\%]}$, %N) defined as g of N per 100 g of dry leaf matter. Each leaf sample was constituted by the leaves of at least three
130 different trees of the dominant tree species in the canopy. The species dominance was determined by the tallest individuals.
The leaves were collected from the upper central part of the crown using extensible loppers. All foliar cohorts present in the
canopy were included in the leaf sample.

The samples were dried and then ground using a Braun Mikrodismembrator-U (B. Braun Biotech International, Melsungen,
Germany). They were analysed for foliar N concentration using the combustion technique coupled to gas chromatography
135 using a Thermo Electron Gas Chromatograph (model NA 2100, CE Instruments-Thermo Electron, Milan, Italy) (Gracia et al.,
2004).

Along with the canopy $N_{[\%]}$ data, we used foliar biomass data (g m^{-2}) acquired during the same forest inventory ($n = 2286$).
The foliar biomass data were obtained for each plot from allometric equations relating the diameter of the branches to the
leaves dry weight (Gracia et al. (2004), Fig. A1). These foliar biomass data were used to calculate canopy N content ($N_{[area]}$, g
140 m^{-2}) for each plot following Eq. (1):

$$\text{canopy } N_{[area]} = \frac{\text{canopy } N_{[\%]} * f_{biom}}{100}, \quad (1)$$

where $\text{canopy } N_{[area]}$ is the canopy N content (N g per square meter of ground area, g m^{-2}), $\text{canopy } N_{[\%]}$ is the canopy N
concentration (%N) and f_{biom} is the foliar biomass (N g per square meter of ground area, g m^{-2}).

Catalonian forests include both deciduous and evergreen broadleaf as well as evergreen needleleaf tree species. These three
145 PFTs are referred to as Deciduous Broadleaf Forest (DBF), Evergreen Broadleaf Forest (EBF) and Evergreen Needleleaf Forest
(ENF), respectively. The main tree species are *Pinus halepensis* Mill., *Pinus sylvestris* L., *Quercus ilex* L., *Pinus uncinata*
Ramond ex DC., *Pinus nigra* J.F. Arnold, *Quercus suber* L., *Quercus cerrioides* Willk. & Costa., *Quercus petraea* Liebl. and
Fagus sylvatica L. These species accounted for 92% of the sampled forest plots. The 15 tree species included in this analysis
are listed in Table 1. Plots with a rare dominant tree species, i.e. species that were detected in only one single plot, were
150 excluded from the analysis. This applied to plots with these dominant species: *Abies alba* Mill., *Fraxinus augustifolia* Vahl,
Fraxinus excelsior L., *Pinus radiata* D. Don, *Populus nigra* L., *Populus tremula* L., *Quercus robur* L.

A proportion of 96% of the plots included in this analysis had a single dominant tree species. The plots with codominant
species ($n = 30$) were sampled several (up to two) times, each time for each of the codominant tree species found on the plots.
After separate measurements, the obtained foliar N concentration values were averaged in order to obtain a single canopy $N_{[\%]}$
155 value for each plot with several codominant species. Among these 30 plots with codominant species, 16 plots had codominant
species from different PFT. Their PFT is thus labelled as mixed while the plots with several codominant species from the same
PFT are labelled according to their PFTs.



2.2.2 MTCI product

160 The MERIS Terrestrial Chlorophyll Index (MTCI) was first developed to estimate chlorophyll content in canopies. Its calculation, presented in Eq. (2), is based on three reflectance bands, located around the red edge point (REP) (Dash and Curran, 2004):

$$MTCI = \frac{R_{band10} - R_{band9}}{R_{band9} - R_{band8}} = \frac{R_{753.75} - R_{708.75}}{R_{708.75} - R_{681.25}} \quad (2)$$

165 where R_{band8} , R_{band9} and R_{band10} represent the 8th, 9th and 10th bands of MERIS, respectively. Following MERIS standard bands settings, the centres of the bands were located at 681.25 nm, 708.75 nm and 753.75 nm on the electromagnetic spectrum. While the ESA ENVISAT satellite mission producing MERIS data came to an end in 2012, MERIS products and MTCI in particular are still relevant because the new ESA Sentinel-2 satellite mission has improved band settings compared to those of MERIS and increased the spatial resolution to 20 m (Drusch et al., 2012). The Sentinel-2 mission will also release a chlorophyll product that will continue the time series already available for MTCI. In this study, we put emphasis on ENVISAT-MERIS as
170 our field data are closer to the MERIS acquisition period.

MTCI has been described to be sensitive to high chlorophyll content while presenting low sensitivity to soil brightness. MTCI 10 year time series are available almost globally and are already corrected for atmospheric influences and cloud cover (Curran and Dash, 2005).

175 MTCI level 3 imagery was obtained from the NERC Earth Observation Data Centre (NEODC, 2015) for the region of Catalonia between 2002 and 2012. The original data were provided by the European Space Agency and then processed by Airbus Defence and Space. The original data, following ENVISAT specifications, have a revisit time of three days and the processed imagery is available as an either weekly or monthly averaged product. The spatial resolution of the processed data is approximately 1 km. One MTCI imagery product covering the entire study area was obtained for every month between June 2002 and March 2012, except for October 2003, when no valid product was available.

180 2.3 Data handling

2.3.1 Methodology to link canopy N data to MTCI values

185 There is a discrepancy between the timing of the ground truth sampling and the satellite image acquisition period. While the plot sampling campaigns were carried out between 1988 and 2001, the ENVISAT satellite mission was launched in 2002 and ended in 2012. To overcome the discrepancy, MTCI images were averaged by month over the 10 years of the satellite mission period. This process yielded twelve MTCI averaged images, one for each month. The averaged MTCI images were then linked to the forest plots based on the forest plot coordinates and sampling month, as the exact sampling date was known for each plot. The period between the 1st of June and the 31st of October was determined to be the growing season after a pre-analysis, where we studied yearly temporal variation of MTCI in several locations and forest types in Catalonia. This extended period was chosen to encompass the different vegetation phenology types corresponding to the contrasted climate conditions in this



190 region. The forest plots sampled outside of the growing season were excluded from the analysis. The Globcover 2009 land
cover map was used to exclude forest plots located on unsuitable land surface. The Globcover map was created by ESA using
MERIS reflectance data from 2009 (Bontemps et al., 2011). It was downloaded from the ESA data user elements website
(ESA, 2010). This map comprises 22 land cover classes and has a spatial resolution of 300 m. Using this map, we excluded
195 forest plots that had undergone a land cover change since the sampling period and did not have a natural vegetation cover any
more at the time of remote sensing image acquisition. To do so, the landcover map was first resampled to a spatial resolution
of 1 km to be in accordance with MTCI spatial resolution. Then, the plots located on land area classified as either rainfed
cropland, mosaic between croplands and natural vegetation, sparse vegetation or artificial surfaces were excluded from the
analysis.

2.3.2 Relationship between MTCI and canopy N data at lower spatial resolution

200 In a first step, the relationships between MTCI and canopy N data values were investigated after resampling both datasets to
the same, lower, spatial resolution. This was done because of the initial difference in support size between MTCI spatial
resolution and the forest plots size (i.e. 1 km and 6 m, respectively). This enabled us to investigate the relationships between
MTCI and canopy N data independently of differences in initial support size.

The Globcover 2009 land cover map was used to exclude from the computation the MTCI pixels located on land surface
205 without natural vegetation cover. As for the forest plots, MTCI pixels whose land cover class corresponded to rainfed cropland,
mosaic between croplands and natural vegetation, sparse vegetation or artificial surfaces were excluded from the upscaling
analysis. The monthly averaged MTCI images obtained previously were resampled successively to 5 km, 10 km, 15 km, and
20 km. Forest plots data were then averaged over the newly obtained pixel. The relationship between the resampled MTCI
values and canopy N data was analysed using linear regression.

2.3.3 Relationship between MTCI and canopy N data at initial higher spatial resolution

210 In a second step, the relationships between MTCI and canopy N data, both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$, were examined at
the original spatial resolution of 1 km. This allowed us to investigate the influence of PFT and species on the relationships as
this information was lost in the resampling process. The relationships between MTCI and canopy N at 1 km-spatial resolution
were analysed with linear regression for the whole dataset, for each PFT separately as well as for individual species.

2.3.4 Statistical analysis

215 After applying the selection criteria explained above, i.e. plots measured between June 1st and October 31st, exclusion of plots
with infrequent species and selection based on Globcover 2009, 846 forest plots were available for analysis, including 841
plots with foliar biomass and canopy N content information. Descriptive statistics of canopy $N_{[\%]}$, foliar biomass and canopy
 $N_{[\text{area}]}$ were then produced for each of the tree species and PFT included in the analysis. The linear regressions between MTCI
220 and canopy N were then performed for both resampled and non-resampled datasets. Preliminary analysis showed that using a



natural logarithm transformation (log) of the canopy N variables was necessary to fulfil linear regression model assumptions, namely normality and homogeneity of variance of the residuals. The minimum number of data points needed to carry out the linear regression analysis was fixed at 10. The spatial analyses were done with the PCRaster software (Karssenberget al., 2010). The statistical analyses were performed in the R environment (R Development Core Team, 2014) and the ggplot2 package was used for the graphics (Wickham, 2009).

3 Results

3.1 Descriptive statistics

Descriptive statistical analysis of canopy $N_{[\%]}$, canopy $N_{[\text{area}]}$ and foliar biomass were performed for each tree species included in the dataset (Table 1). The four most abundant species (*Pinus halepensis*, *Pinus sylvestris*, *Quercus ilex* and *Pinus uncinata*) dominated 667 plots i.e. almost 80% of the plots. The cumulated abundance percentages of ENF, EBF and DBF species were equal to 66 %, 22 % and 9 %, respectively. From this data, it is clear that the forests plots were mainly dominated by ENF species. On average, *Pinus uncinata* plots had the highest biomass values while *Quercus suber* plots showed the lowest mean value for this variable. Descriptive statistics were also analysed by PFT. The mean canopy $N_{[\%]}$ was lowest for ENF species, 0.97 %N, and highest for DBF trees, 2.17 %N (Fig. 2a). Canopy $N_{[\%]}$ value ranges were equal to 1.91 %N, 2.06 %N, 1.68 %N and 1.42 %N for DBF, EBF, ENF and mixed plots, respectively. The canopy $N_{[\text{area}]}$ statistics were analysed by PFT as well (Fig. 2b) and the averaged canopy $N_{[\text{area}]}$ values ranged from 1.82 g m⁻² to 4.61 g m⁻². A Pearson correlation matrix (Fig. 3) was computed between the variables for the whole dataset and while the correlation between each pair of variables was significant, the relationship between canopy $N_{[\text{area}]}$ and foliar biomass was strongest ($r = 0.88$). This matrix also shows distribution histograms of the three variables. As canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ distributions are positively skewed, a logarithmic transformation was applied to these variables to fulfil linear model assumptions. Correlation matrices for each DBF, EBF and ENF plots are presented in the Appendix (Fig. A2 – 4).

3.2 Relationship between MTCI and canopy N data at lower spatial resolution

The relationships between MTCI and both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ were studied after resampling both datasets to the same, lower, spatial resolution. This was done to investigate the relationship between MTCI and canopy N data independently of differences in support size. The results showed that the relationship between MTCI and canopy $N_{[\%]}$ was always stronger than the relationship for MTCI and canopy $N_{[\text{area}]}$ for each resampling factor. Moreover, the relationships between MTCI and either canopy $N_{[\%]}$ or canopy $N_{[\text{area}]}$ were all highly significant ($p < 0.000$). The r^2 values of the relationship between MTCI and canopy $N_{[\%]}$ were equal to 0.33, 0.37, 0.35 and 0.43 for 5 km, 10 km, 15 km and 20 km resampled spatial resolution, respectively. The r^2 values of the relationship between MTCI and canopy $N_{[\text{area}]}$ were equal to 0.17, 0.20, 0.20 and 0.17 at 5 km, 10 km, 15 km and 20 km spatial resolution. The relationship between MTCI and canopy $N_{[\%]}$ at 20 km spatial resolution is shown in Figure 4.



3.3 Relationship between MTCI and canopy N data at higher spatial resolution

3.3.1 Relationship between MTCI and canopy N concentration

255 The relationships between MTCI and canopy N data were studied after resampling both datasets to the same, lower, spatial resolution. The results showed that the linear regression between MTCI and canopy $N_{[\%]}$ for the whole dataset ($n = 846$) was highly significant ($p < 0.000$) and had a r^2 value of 0.32 (Table 2, Fig. 5a). The relationship between MTCI and Canopy $N_{[\%]}$ was also investigated for each PFT individually (Fig. 5b-e). For DBF plots, the relationship between MTCI and canopy $N_{[\%]}$ had an r^2 value of 0.25 ($n = 80$) and was significant. However, although significant, the r^2 of the relationship between MTCI and canopy $N_{[\%]}$ for EBF and ENF plots were lower and equal to 0.03 ($n = 186$) and 0.10 ($n = 564$), respectively.

260 The relationship between MTCI and canopy $N_{[\%]}$ was also significant for one individual species, *Fagus sylvatica*. The proportion of explained variance for this species was equal to 0.71 ($n = 15$). This result, although obtained on a restricted number of plots, showed that the significant relationships between MTCI and canopy $N_{[\%]}$ not only existed when all DBF plots were included but also held for one individual DBF species.

3.3.2 Relationship between MTCI and canopy N content

265 Significant relationships between MTCI and canopy $N_{[\text{area}]}$ were found for the whole dataset as for EBF and ENF plots (Table 3). The scatterplots between MTCI and canopy $N_{[\text{area}]}$ are presented in Figure 6. The proportion of explained variance was higher for ENF plots compared to the other PFTs and compared to the overall relationship across all plots. The relationship between MTCI and canopy $N_{[\text{area}]}$ was also investigated for 10 individual species and one of them showed significant relationships: *Quercus ilex*.

270 4 Discussion

Our aim was to explore the relationship between the MTCI vegetation index and both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ in Mediterranean forests at the regional scale in Catalonia, north eastern of Spain. This was done by using the ESA spaceborne MTCI remote sensing product and canopy N data from a forest inventory. The relationship was first investigated using MTCI and canopy N data resampled to the same, lower, spatial resolution. The relationship was then investigated across all plots and
275 by PFT at MTCI initial spatial resolution of 1 km.

4.1 Relationship between MTCI and canopy N data at lower spatial resolution

This pre-analysis was undertaken to study the MTCI-canopy N relationships independently of the discrepancy between MTCI original spatial resolution (1 km) and the size of the forest plots (diameter of 6 m). By resampling both datasets to a lower spatial resolution, the obtained values were less impacted by small-scale variations because they were obtained by averaging
280 several values over a larger area. The results showed that the relationship between MTCI and canopy N data was significant



and consistent across all spatial resolutions investigated: 5 km, 10 km, 15 km and 20 km. This shows that, when the influence of the discrepancy between the original datasets was taken into account, MTCI and canopy N data were linked and that the MTCI-canopy N relationship was not strongly affected by the resampled spatial resolution.

4.2 Relationship between MTCI and canopy N data at higher spatial resolution

285 4.2.1 Canopy N concentration detection

The overall relationship between MTCI and canopy $N_{[\%]}$ at 1 km spatial resolution for all the forest plots ($n = 846$) was significant and the r^2 value was equal to 0.32 (Fig. 5). This result showed that canopy $N_{[\%]}$ could be related to MTCI in Mediterranean forests. The performance of the MTCI vegetation index to detect canopy $N_{[\%]}$ in Mediterranean vegetation was similar to the results obtained from previous studies using spaceborne MTCI. For example, using MTCI computed from the spaceborne RapidEye sensor at 5 m spatial resolution, it was possible to detect canopy $N_{[\%]}$ in grassland savannah and subtropical forest with similar coefficients of determination, $r^2 = 0.35$ and $r^2 = 0.52$, respectively (Ramoelo et al., 2012; Cho et al., 2013). More generally, while there is a consensus regarding MTCI ability for in situ leaf or canopy $N_{[\%]}$ detection in a variety of crops (Tian et al., 2011; Li et al., 2014), there is no general agreement about MTCI ability for canopy $N_{[\%]}$ detection across vegetation and sensor types. For example, MTCI computed from airborne data at 3 m spatial resolution could not be related to canopy $N_{[\%]}$ from a mixed temperate forest (Wang et al., 2016). In this context our finding brings new insight into MTCI $N_{[\%]}$ sensing capabilities at a much coarser spatial resolution (1 km) compared to what has been done before.

Investigating the influence of the PFTS on the overall relationship highlighted the difference between DBF, EBF and ENF types of vegetation regarding canopy $N_{[\%]}$ detection by spaceborne MTCI. The relationships between MTCI and canopy $N_{[\%]}$ were significant for all the PFT taken separately (p -value < 0.05). However, a higher proportion of variance was explained for DBF and mixed plots ($r^2 = 0.25$ and $r^2 = 0.47$ for DBF and mixed plots, respectively) compared to the other plant functional types ($r^2 = 0.10$ and $r^2 = 0.03$ for ENF and EBF trees, respectively) and the relationship between MTCI and canopy $N_{[\%]}$ was especially weaker for EBF plots. Therefore, the relationship observed for all the forest plots may be mainly driven by DBF and mixed plots. This result is different from what was observed by Ollinger et al. (2008) in boreal forests, where canopy $N_{[\%]}$ was related to NIR reflectance for both broadleaf and needleleaf plots taken separately. Moreover, the results obtained for ENF tree species are surprising as previous studies investigating the relationship between foliar $N_{[\%]}$ and in situ measured spectra reported higher r^2 values, $r^2 = 0.59$ and $r^2 = 0.81$ in spruce and pine forest, respectively (Stein et al., 2014; Schlerf et al., 2010). The differences in scale and methodology might explain the divergent results compared to previous findings. Indeed, in our study, the analysis is carried out at a much coarser spatial resolution using spaceborne data compared to the fine spatial scale obtained with in situ devices. Moreover, most of these studies were carried out in temperate forests and studies investigating canopy $N_{[\%]}$ detection in Mediterranean regions are scarce. When investigating the relationship between canopy $N_{[\%]}$ and MTCI at the species level, we also found that it was significant for *Fagus sylvatica* plots ($r^2 = 0.71$).



315 In the literature, the relationship between MTCI and canopy $N_{[\%]}$ is often not stratified by PFT or species (Sullivan et al., 2013; Wang et al., 2016). In this study, we showed that investigating this relationship for each PFT taken separately yielded additional insight. Indeed, to our knowledge the difference in explained variance between DBF and other PFTs in MTCI and canopy $N_{[\%]}$ relationship has not been observed before. Moreover, we could show that the stronger relationship observed for DBF plots was consistent for *Fagus sylvatica* plots taken separately.

4.2.2 Canopy N content detection

320 The relationship between MTCI and canopy $N_{[\text{area}]}$, which was obtained by combining canopy N concentration values with biomass data, was significant across all plots ($n = 841$) (Table 3, Fig. 6). Although the r^2 value was lower for the relationship between MTCI and canopy $N_{[\text{area}]}$ ($r^2 = 0.17$) than for the relationship between MTCI and canopy $N_{[\%]}$ ($r^2 = 0.32$), it is interesting to note that canopy $N_{[\text{area}]}$ can be related to spaceborne MTCI as remotely sensed detection of canopy $N_{[\text{area}]}$ is rarely investigated in forest environments (Mirik et al., 2005). In comparison, previous studies conducted in grasslands reported higher prediction accuracy e.g. by using spaceborne MTCI at 300 m spatial resolution or a simple ratio-type vegetation index computed from airborne imagery at 1 m spatial resolution, canopy $N_{[\text{area}]}$ was detected with r^2 values equal to 0.29 and 0.66, respectively (Mirik et al., 2005; Ullah et al., 2012).

325 The relationship between MTCI and canopy $N_{[\text{area}]}$ was only significant for ENF and EBF plots (Fig. 6b-e), with a higher proportion of explained variance for ENF plots ($r^2 = 0.20$). However, when this relationship was investigated at the species scale, significant results were found for *Quercus ilex* (EBF) plots. This is accordance with a previous study examining the detection of canopy $N_{[\text{area}]}$ in *Quercus ilex* trees by MTCI computed from in situ spectra ($r^2 = 0.43$) (Pacheco-Labrador et al., 330 2014).

4.3 Comparing results obtained for canopy N concentration and canopy N content detection

This analysis highlighted the difference between canopy N expressed as a percentage of leaf dry matter (canopy $N_{[\%]}$) and on an area basis (canopy $N_{[\text{area}]}$) regarding detection by spaceborne MTCI for the different PFTs. Canopy $N_{[\%]}$ of DBF and mixed plots showed higher correlation with MTCI compared to EBF and ENF plots while the relationship between canopy $N_{[\text{area}]}$ of ENF plots with MTCI was stronger than for any other PFTs. These differences between canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ detection by remote sensing can be related to previous findings showing that canopy $N_{[\text{area}]}$ but not canopy $N_{[\%]}$ could be detected by MTCI in grassland (Ullah et al., 2012) and by a simple ratio index in heterogeneous rangelands (Mirik et al., 2005) at various spatial scales, 300 m and 1 m, respectively. In the literature, canopy $N_{[\%]}$ is more often used to detect N state of foliage in forest while canopy $N_{[\text{area}]}$ is regularly employed in grasslands but also in crops (Clevers and Gitelson, 340 2013; Schlemmer et al., 2013). Our results showed that, for ENF plots, when biomass was accounted for, as in canopy $N_{[\text{area}]}$, the relationship between MTCI and canopy $N_{[\text{area}]}$ was stronger compared to canopy $N_{[\%]}$.



4.4 Influence of the plant functional type on the MTCI canopy N relationship

The relationships between MTCI and both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ were influenced by the PFT of the plots. The relationship between MTCI and canopy $N_{[\%]}$ was stronger for DBF and mixed plots compared to EBF and ENF plots while the opposite was true for the MTCI-canopy $N_{[\text{area}]}$ relationship. In the ongoing discussion about the mechanisms underlying the remote detection of canopy N, some authors argued that the differences in canopy structure between different PFTs were causing the observed relationship between canopy N and remote sensing data, rendering it spurious (Knyazikhin et al., 2013). Other authors, although agreeing that canopy structural properties needed to be accounted for, suggested that a direct biochemical link between canopy N and reflectance data was not necessary to detect canopy N with reflectance data (Ollinger et al., 2013; Townsend et al., 2013). In this context, our analysis showed that the PFTs of the plots had an influence on the MTCI canopy N relationship in a specific type of ecosystem, namely Mediterranean forests. Further analysis is needed to disentangle the influence of structural properties on canopy N detection.

4.5 Perspectives for larger scale applications

The methodology applied in this paper is different from the usual methodology implemented to detect canopy N concentration in forests. Remote sensing of N in forest canopies by hyperspectral sensors is often coupled with intensive forest sampling measurements. This method has been effective at detecting canopy N concentration locally in a vast range of environments (Serrano et al., 2002; Smith et al., 2002; Townsend et al., 2003; Ollinger et al., 2008; Wang et al., 2016). Applying this technique at larger scales has already been explored. For example, Martin *et al.*, (2008) compiled 137 field plots data from previous studies in various forest types and investigated the possibility to find a common detection algorithm. However, due to the different treatments required as well as the high spatial resolution (from 3 m to 30 m for Hypex airborne and Hyperion spaceborne sensors, respectively, Wang et al., 2016; Smith et al., 2003), applying imaging spectroscopy at a broader scale might reveal laborious. A recent study in northern temperate forests explored the effect of spatial resolution on canopy $N_{[\%]}$ estimation. The results showed that, although the prediction accuracy was reduced compared to what was achieved using PLS regression at higher spatial resolution, it was still possible to estimate canopy $N_{[\%]}$ with r^2 between 0.34 and 0.81 using various vegetation indices computed from MODIS reflectance data at 500 m spatial resolution (Lepine et al., 2016). In this context, the methodology applied in this article could be a valuable alternative to explore canopy N detection at larger scale. Using published data from an extensive field plot inventory, we were able to relate both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ to MTCI at different spatial resolutions. As MTCI time series are readily and almost globally available, it could eventually be possible to apply our approach at a broader scale in different types of biomes. The results obtained for DBF species and *Fagus sylvatica* in particular suggest that this method may be efficient at estimating canopy N in temperate forests. Obtaining reliable canopy N estimates over larger scale and for diverse and globally distributed type of vegetation would be necessary to be able to use such estimates to improve and calibrate global vegetation models.



5 Conclusion

In this study, we investigated the relationship between spaceborne MTCI from ENVISAT and both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$ at regional scale in Mediterranean forests. We found significant results across all plots both when the original data were resampled to 5 km, 10 km, 15 km and 20 km and for the original spatial resolution of 1 km. The relationship between MTCI and canopy N data was also significant for some individual PFTs and species. The r^2 values were 0.32 and 0.17 for the overall relationships between MTCI and either canopy $N_{[\%]}$ or canopy $N_{[\text{area}]}$, respectively. We highlighted the differences between PFTs and both canopy $N_{[\%]}$ and canopy $N_{[\text{area}]}$: the relationship between MTCI and canopy $N_{[\%]}$ was stronger for DBF and mixed plots while canopy $N_{[\text{area}]}$ was more linked to MTCI for ENF plots. Such differences in relationships between MTCI and either canopy $N_{[\%]}$ or canopy $N_{[\text{area}]}$ were already observed in grasslands ecosystem. Our results showed that MTCI could be related to canopy N for some individual PFTs, indicating an influence of the PFTs on the MTCI-canopy N relationship. The methodology developed in this study could be investigated at larger scales in different types of ecosystem. While this could already be undertaken using the ENVISAT MTCI 10 years time series as it is almost globally available, ESA new Sentinel-2 satellite launched on 23 June 2015 yields reflectance data at improved spatial and temporal resolution than ENVISAT-MERIS. Canopy N estimates collected through larger scales applications could be exploited in vegetation modelling studies including both the C and N cycles.

6 Data availability

The canopy data used in this study can be obtained from the TRY Plant Trait Database (<https://www.try-db.org/TryWeb/Home.php>, dataset 91) or by directly contacting the authors.

7 Appendix A

This appendix presents the Allometric relationship between the branch diameter (mm) and the foliar dry mass (g) from which the biomass data was obtained (Fig. A1) and the correlation matrices for DBF (Fig. A2), EBF (Fig. A3) and ENF plots (Fig. A3).

8 Competing interest

The authors declare that they have no conflict of interest.



9 Acknowledgements

This research was funded by The Netherlands Organisation for Scientific Research (NWO) under the project number NWO ALW-GO-AO/14-12. J. Sardans and J. Peñuelas were funded by the European Research Council Synergy grant SyG-2013-610028 IMBALANCE-P, the Spanish Government projects CGL2013-48074-P and the Catalan Government project SGR 2014-274. We would like to acknowledge Scott Ollinger and Lucie Lepine for their valuable comments and discussions on our research project as well as Ton Markus for his help with the figures presented in this paper.

405



10 References

410 Boyd, D. S., Almond, S., Dash, J., Curran, P. J., and Hill, R. A.: Phenology of vegetation in southern England from Envisat
MERIS terrestrial chlorophyll index (MTCI) data, *International Journal of Remote Sensing*, 32, 8421-8447,
doi:10.1080/01431161.2010.542194, 2011.

Boyd, D. S., Almond, S., Dash, J., Curran, P. J., Hill, R. A., and Foody, G. M.: Evaluation of Envisat MERIS terrestrial
chlorophyll index-based models for the estimation of terrestrial gross primary productivity, *IEEE Geoscience and Remote
Sensing Letters*, 9, 457-461, 2012.

415 Cho, M. A., Ramoelo, A., Debba, P., Mutanga, O., Mathieu, R., van Deventer, H., and Ndlovu, N.: Assessing the effects
of subtropical forest fragmentation on leaf nitrogen distribution using remote sensing data, *Landscape Ecology*, 28, 1479-1491,
doi:10.1007/s10980-013-9908-7, 2013.

Clevers, J. G. P. W., and Gitelson, A. A.: Remote estimation of crop and grass chlorophyll and nitrogen content using
red-edge bands on Sentinel-2 and -3, *International Journal of Applied Earth Observation and Geoinformation*, 23, 344-351,
doi:10.1016/j.jag.2012.10.008, 2013.

420 Crabbe, R. A., Dash, J., Rodriguez-Galiano, V. F., Janous, D., Pavelka, M., and Marek, M. V.: Extreme warm
temperatures alter forest phenology and productivity in Europe, *Science of The Total Environment*, 563-564, 486-495,
<http://dx.doi.org/10.1016/j.scitotenv.2016.04.124>, 2016.

Curran, P. J., and Dash, J.: Algorithm Theoretical Basis Document ATBD 2.22 Chlorophyll Index, University of
Southampton, Southampton, 41 pp., available at:
425 https://earth.esa.int/documents/700255/2042855/MERIS_ATBD_2042852.2042822_v2042851.2042852++2042005.pdf,
last access 2042811 November 2042016, 2005.

Dash, J., and Curran, P. J.: The MERIS terrestrial chlorophyll index, *International Journal of Remote Sensing*, 25, 5403-
5413, doi:10.1080/0143116042000274015, 2004.

430 Dash, J., and Curran, P. J.: Evaluation of the MERIS terrestrial chlorophyll index (MTCI), *Advances in Space Research*,
39, 100-104, doi:10.1016/j.asr.2006.02.034, 2007.

Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P.,
Martimort, P., Meygret, A., Spoto, F., Sy, O., Marchese, F., and Bargellini, P.: Sentinel-2: ESA's Optical High-Resolution
Mission for GMES Operational Services, *Remote Sensing of Environment*, 120, 25-36, doi:10.1016/j.rse.2011.11.026, 2012.

435 Evans, J. R.: Photosynthesis and nitrogen relationships in leaves of C3 plants, *Oecologia*, 78, 9-19,
doi:10.1007/BF00377192, 1989.

Ferwerda, J. G., Skidmore, A. K., and Mutanga, O.: Nitrogen detection with hyperspectral normalized ratio indices across
multiple plant species, *International Journal of Remote Sensing*, 26, 4083-4095, 10.1080/01431160500181044, 2005.

Filella, I., and Penuelas, J.: The red edge position and shape as indicators of plant chlorophyll content, biomass and hydric
status, *International Journal of Remote Sensing*, 15, 1459-1470, <http://dx.doi.org/10.1080/01431169408954177>, 1994.



- 440 Filella, I., Serrano, L., Serra, J., and Penuelas, J.: Evaluating wheat nitrogen status with canopy reflectance indices and discriminant analysis, *Crop Science*, 35, 1400-1405, doi:10.2135/cropsci1995.0011183X003500050023x, 1995.
- Foody, G. M., and Dash, J.: Discriminating and mapping the C3 and C4 composition of grasslands in the northern Great Plains, USA, *Ecological Informatics*, 2, 89-93, doi:10.1016/j.ecoinf.2007.03.009, 2007.
- 445 Glenn, E. P., Huete, A. R., Nagler, P. L., and Nelson, S. G.: Relationship between remotely-sensed vegetation indices, canopy attributes and plant physiological processes: What vegetation indices can and cannot tell us about the landscape, *Sensors*, 8, 2136-2160, doi:10.3390/s8042136, 2008.
- González, J. R., and Pukkala, T.: Characterization of forest fires in Catalonia (north-east Spain), *European Journal of Forest Research*, 126, 421-429, doi:10.1007/s10342-006-0164-0, 2007.
- 450 Gracia, C., Ibàñez, J. J., Burriel, J. A., Mata, T., and Vayreda, J.: Inventari Ecològic i Forestal de Catalunya. Mètodes, CREAM, Bellaterra, available at: <http://www.creaf.uab.es/iefc/pub/Metodes/index.htm>, last access 11 November 2016, 2004.
- Green, D. S., Erickson, J. E., and Kruger, E. L.: Foliar morphology and canopy nitrogen as predictors of light-use efficiency in terrestrial vegetation, *Agricultural and Forest Meteorology*, 115, 163-171, [http://dx.doi.org/10.1016/S0168-1923\(02\)00210-1](http://dx.doi.org/10.1016/S0168-1923(02)00210-1), 2003.
- 455 Hansen, P. M., and Schjoerring, J. K.: Reflectance measurement of canopy biomass and nitrogen status in wheat crops using normalized difference vegetation indices and partial least squares regression, *Remote Sensing of Environment*, 86, 542-553, doi:10.1016/S0034-4257(03)00131-7, 2003.
- Harris, A., and Dash, J.: The potential of the MERIS Terrestrial Chlorophyll Index for carbon flux estimation, *Remote Sensing of Environment*, 114, 1856-1862, doi:10.1016/j.rse.2010.03.010, 2010.
- 460 Harris, A., and Dash, J.: A new approach for estimating northern peatland gross primary productivity using a satellite-sensor-derived chlorophyll index, *Journal of Geophysical Research: Biogeosciences*, 116, doi:10.1029/2011JG001662, 2011.
- Huber, S., Kneubühler, M., Psomas, A., Itten, K., and Zimmermann, N. E.: Estimating foliar biochemistry from hyperspectral data in mixed forest canopy, *Forest Ecology and Management*, 256, 491-501, doi:10.1016/j.foreco.2008.05.011, 2008.
- 465 Kergoat, L., Lafont, S., Arneth, A., Le Dantec, V., and Saugier, B.: Nitrogen controls plant canopy light-use efficiency in temperate and boreal ecosystems, *Journal of Geophysical Research: Biogeosciences*, 113, doi:10.1029/2007JG000676, 2008.
- 470 Knyazikhin, Y., Schull, M. A., Stenberg, P., Möttus, M., Rautiainen, M., Yang, Y., Marshak, A., Latorre Carmona, P., Kaufmann, R. K., Lewis, P., Disney, M. I., Vanderbilt, V., Davis, A. B., Baret, F., Jacquemoud, S., Lyapustin, A., and Myneni, R. B.: Hyperspectral remote sensing of foliar nitrogen content, *Proceedings of the National Academy of Sciences*, 110, E185–E192, [10.1073/pnas.1210196109](https://doi.org/10.1073/pnas.1210196109), 2013.
- Lana, X., and Burgueño, A.: Spatial and temporal characterization of annual extreme droughts in Catalonia (Northeast Spain), *International Journal of Climatology*, 18, 93-110, 1998.



Lepine, L. C., Ollinger, S. V., Ouimette, A. P., and Martin, M. E.: Examining spectral reflectance features related to foliar nitrogen in forests: Implications for broad-scale nitrogen mapping, *Remote Sensing of Environment*, 173, 174-186, doi:10.1016/j.rse.2015.11.028, 2016.

Li, F., Miao, Y., Feng, G., Yuan, F., Yue, S., Gao, X., Liu, Y., Liu, B., Ustin, S. L., and Chen, X.: Improving estimation of summer maize nitrogen status with red edge-based spectral vegetation indices, *Field Crops Research*, 157, 111-123, doi:10.1016/j.fcr.2013.12.018, 2014.

Liu, D., Ogaya, R., Barbeta, A., Yang, X., and Peñuelas, J.: Contrasting impacts of continuous moderate drought and episodic severe droughts on the aboveground-biomass increment and litterfall of three coexisting Mediterranean woody species, *Global Change Biology*, 21, 4196-4209, doi:10.1111/gcb.13029, 2015.

Martin, M. E., Plourde, L. C., Ollinger, S. V., Smith, M. L., and McNeil, B. E.: A generalizable method for remote sensing of canopy nitrogen across a wide range of forest ecosystems, *Remote Sensing of Environment*, 112, 3511-3519, doi:10.1016/j.rse.2008.04.008, 2008.

Mirik, M., Norland, J. E., Crabtree, R. L., and Biondini, M. E.: Hyperspectral one-meter-resolution remote sensing in Yellowstone National Park, Wyoming: I. Forage nutritional values, *Rangeland Ecology and Management*, 58, 452-458, doi:10.2111/04-17.1, 2005.

NEODC - NERC Earth Observation Data Centre: <http://neodc.nerc.ac.uk/>, access: 06/02/2015, 2015.

Ollinger, S. V., and Smith, M. L.: Net primary production and canopy nitrogen in a temperate forest landscape: An analysis using imaging spectroscopy, modeling and field data, *Ecosystems*, 8, 760-778, doi:10.1007/s10021-005-0079-5, 2005.

Ollinger, S. V., Richardson, A. D., Martin, M. E., Hollinger, D. Y., Frohling, S. E., Reich, P. B., Plourde, L. C., Katul, G. G., Munger, J. W., Oren, R., Smith, M. L., Paw U, K. T., Bolsta, P. V., Cook, B. D., Day, M. C., Martin, T. A., Monson, R. K., and Schmid, H. P.: Canopy nitrogen, carbon assimilation, and albedo in temperate and boreal forests: Functional relations and potential climate feedbacks, *Proceedings of the National Academy of Sciences of the United States of America*, 105, 19336-19341, doi:10.1073/pnas.0810021105., 2008.

Ollinger, S. V., Reich, P. B., Frohling, S., Lepine, L. C., Hollinger, D. Y., and Richardson, A. D.: Nitrogen cycling, forest canopy reflectance, and emergent properties of ecosystems, *Proceedings of the National Academy of Sciences*, 110, E2437, 10.1073/pnas.1304176110, 2013.

Pacheco-Labrador, J., González-Cascón, R., Pilar Martín, M., and Riaño, D.: Understanding the optical responses of leaf nitrogen in mediterranean holm oak (*Quercus ilex*) using field spectroscopy, *International Journal of Applied Earth Observation and Geoinformation*, 26, 105-118, doi:10.1016/j.jag.2013.05.013, 2014.

Peng, Y., and Gitelson, A. A.: Application of chlorophyll-related vegetation indices for remote estimation of maize productivity, *Agricultural and Forest Meteorology*, 151, 1267-1276, doi:10.1016/j.agrformet.2011.05.005, 2011.

Peñuelas, J., Gamon, J. A., Fredeen, A. L., Merino, J., and Field, C. B.: Reflectance indices associated with physiological changes in nitrogen- and water-limited sunflower leaves, *Remote Sensing of Environment*, 48, 135-146, doi:10.1016/0034-4257(94)90136-8, 1994.



- Ramoelo, A., Skidmore, A. K., Cho, M. A., Schlerf, M., Mathieu, R., and Heitkönig, I. M. A.: Regional estimation of savanna grass nitrogen using the red-edge band of the spaceborne rapideye sensor, *International Journal of Applied Earth Observation and Geoinformation*, 19, 151-162, doi:10.1016/j.jag.2012.05.009, 2012.
- 510 Reich, P. B., Walters, M. B., Kloeppel, B. D., and Ellsworth, D. S.: Different photosynthesis-nitrogen relations in deciduous hardwood and evergreen coniferous tree species, *Oecologia*, 104, 24-30, doi:10.1007/BF00365558, 1995.
- Reich, P. B., Walters, M. B., and Ellsworth, D. S.: From tropics to tundra: Global convergence in plant functioning, *Proceedings of the National Academy of Sciences of the United States of America*, 94, 13730-13734, doi:10.1073/pnas.94.25.13730, 1997.
- 515 Reich, P. B., Ellsworth, D. S., Walters, M. B., Vose, J. M., Gresham, C., Volin, J. C., and Bowman, W. D.: Generality of leaf trait relationships: A test across six biomes, *Ecology*, 80, 1955-1969, doi:10.2307/176671 1999.
- Reich, P. B.: Key canopy traits drive forest productivity, *Proceedings of the Royal Society B: Biological Sciences*, 279, 2128-2134, doi:10.1098/rspb.2011.2270, 2012.
- Rodriguez-Galiano, V. F., Dash, J., and Atkinson, P. M.: Characterising the land surface phenology of Europe using
520 decadal MERIS data, *Remote Sensing*, 7, 9390-9409, doi:10.3390/rs70709390, 2015.
- Sardans, J., Rivas-Ubach, A., and Peñuelas, J.: Factors affecting nutrient concentration and stoichiometry of forest trees in Catalonia (NE Spain), *Forest Ecology and Management*, 262, 2024-2034, doi:10.1016/j.foreco.2011.08.019, 2011.
- Schlemmer, M., Gitelson, A., Schepers, J., Ferguson, R., Peng, Y., Shanahan, J., and Rundquist, D.: Remote estimation of nitrogen and chlorophyll contents in maize at leaf and canopy levels, *International Journal of Applied Earth Observation and Geoinformation*, 25, 47-54, doi:10.1016/j.jag.2013.04.003, 2013.
- 525 Schlerf, M., Atzberger, C., Hill, J., Buddenbaum, H., Werner, W., and Schüler, G.: Retrieval of chlorophyll and nitrogen in Norway spruce (*Picea abies* L. Karst.) using imaging spectroscopy, *International Journal of Applied Earth Observation and Geoinformation*, 12, 17-26, doi:10.1016/j.jag.2009.08.006, 2010.
- Serrano, L., Peñuelas, J., and Ustin, S. L.: Remote sensing of nitrogen and lignin in Mediterranean vegetation from
530 AVIRIS data: Decomposing biochemical from structural signals, *Remote Sensing of Environment*, 81, 355-364, doi:10.1016/S0034-4257(02)00011-1, 2002.
- Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model, *Biogeosciences*, 11, 2027-2054, doi:10.5194/bg-11-2027-2014, 2014.
- 535 Smith, M. L., Ollinger, S. V., Martin, M. E., Aber, J. D., Hallett, R. A., and Goodale, C. L.: Direct estimation of aboveground forest productivity through hyperspectral remote sensing of canopy nitrogen, *Ecological Applications*, 12, 1286-1302, doi:10.2307/3099972 2002.
- Smith, M. L., Martin, M. E., Plourde, L., and Ollinger, S. V.: Analysis of hyperspectral data for estimation of temperate forest canopy nitrogen concentration: Comparison between an airborne (AVIRIS) and a spaceborne (Hyperion) sensor, *IEEE Transactions on Geoscience and Remote Sensing*, 41, 1332-1337, doi:10.1109/TGRS.2003.813128, 2003.
- 540



Stein, B. R., Thomas, V. A., Lorentz, L. J., and Strahm, B. D.: Predicting macronutrient concentrations from loblolly pine leaf reflectance across local and regional scales, *GIScience and Remote Sensing*, 51, 269-287, doi:10.1080/15481603.2014.912875, 2014.

545 Sullivan, F. B., Ollinger, S. V., Martin, M. E., Ducey, M. J., Lepine, L. C., and Wicklein, H. F.: Foliar nitrogen in relation to plant traits and reflectance properties of New Hampshire forests, *Canadian Journal of Forest Research*, 43, 18-27, doi:10.1139/cjfr-2012-0324, 2013.

Tian, Y. C., Yao, X., Yang, J., Cao, W. X., Hannaway, D. B., and Zhu, Y.: Assessing newly developed and published vegetation indices for estimating rice leaf nitrogen concentration with ground- and space-based hyperspectral reflectance, *Field Crops Research*, 120, 299-310, doi:10.1016/j.fcr.2010.11.002, 2011.

550 Townsend, P. A., Foster, J. R., Chastain Jr, R. A., and Currie, W. S.: Application of imaging spectroscopy to mapping canopy nitrogen in the forest of the central Appalachian mountains using hyperion and AVIRIS, *IEEE Transactions on Geoscience and Remote Sensing*, 41, 1347-1354, doi:10.1109/TGRS.2003.813205, 2003.

555 Townsend, P. A., Serbin, S. P., Kruger, E. L., and Gamon, J. A.: Disentangling the contribution of biological and physical properties of leaves and canopies in imaging spectroscopy data, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 10.1073/pnas.1300952110, 2013.


Ullah, S., Si, Y., Schlerf, M., Skidmore, A. K., Shafique, M., and Iqbal, I. A.: Estimation of grassland biomass and nitrogen using MERIS data, *International Journal of Applied Earth Observation and Geoinformation*, 19, 196-204, doi:10.1016/j.jag.2012.05.008, 2012.

560 Wang, Z., Wang, T., Darvishzadeh, R., Skidmore, A. K., Jones, S., Suarez, L., Woodgate, W., Heiden, U., Heurich, M., and Hearne, J.: Vegetation indices for mapping canopy foliar nitrogen in a mixed temperate forest, *Remote Sensing*, 8, doi:10.3390/rs8060491, 2016.

565 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C., Midgley, J. J., Navas, M.-L., Niinemets, U., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V. I., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The worldwide leaf economics spectrum, *Nature*, 428, 821-827, doi:10.1038/nature02403, 2004.


Zaehle, S., and Friend, A.: Carbon and nitrogen cycle dynamics in the O - CN land surface model: 1. Model description, site - scale evaluation, and sensitivity to parameter estimates, *Global Biogeochemical Cycles*, 24, doi:10.1029/2009GB003521, 2010.

570



9251

Figure 1. Map showing the forest plots location in the region of Catalonia, north eastern Spain. DBF = Deciduous Broadleaf Forest, EBF = Evergreen Broadleaf Forest, ENF = Evergreen Needleleaf Forest, mixed = mixed forest.



575

Figure 2. Boxplot of (a) canopy nitrogen (N) concentration (canopy $N_{[\%]}$, %N) for Deciduous Broadleaf Forest plots (DBF, $n = 80$), Evergreen Broadleaf Forest plots (EBF, $n = 186$), Evergreen Needleleaf Forest plots (ENF, $n = 564$) and mixed forest plots (mixed, $n = 16$); (b) canopy N content (canopy $N_{[area]}$, $g\ m^{-2}$) for Deciduous Broadleaf Forest plots (DBF, $n = 80$), Evergreen Broadleaf Forest plots (EBF, $n = 186$), Evergreen Needleleaf Forest plots (ENF, $n = 563$) and mixed forest plots (mixed, $n = 12$);

580





Figure 3. The upper right part of this figure shows the Pearson correlation matrix between canopy N_[%] (%N), canopy N_[area] (g m⁻²) and foliar biomass (g m⁻²) variables for the whole dataset, n = 841. The diagonal presents the histograms of the variables on the x-axis, while the y-axis represents the number of counts. The lower left part of this figure represents the scatterplots between the variables. PFT = Plant Functional Type, DBF = Deciduous Broadleaf Forest, EBF = Evergreen Broadleaf Forest, ENF = Evergreen Needleleaf Forest, mixed = mixed forest.

585



590 **Figure 4.** Scatterplot between the MERIS Terrestrial Chlorophyll Index (MTCI) (-) and canopy nitrogen concentration (canopy $N_{[\%]}$, %N) after resampling the datasets to 20 km-spatial resolution ($n = 204$).




Figure 5. Scatterplot and linear regression line between the MERIS Terrestrial Chlorophyll Index (MTCI) (-) and canopy nitrogen (N) concentration (canopy N [%], %N) for (a) whole dataset (n = 846); (b) Deciduous Broadleaf Forest plots (DBF, n = 80); (c) Evergreen Broadleaf Forest plots (EBF, n = 186); (d) Evergreen Needleleaf Forest plots (ENF, n = 564); (e) mixed forest plots (n = 16). PFT = Plant functional type. Canopy N [%] variable was log transformed to fulfil linear model assumptions.

595




Figure 6. Scatterplot and linear regression line between the MERIS Terrestrial Chlorophyll Index (MTCI) (-) and canopy N content ($\text{canopy } N_{[\text{area}]}$, g m^{-2}) for (a) whole dataset ($n = 841$); (b) Deciduous Broadleaf Forest plots (DBF, $n = 80$); (c) Evergreen Broadleaf Forest plots (EBF, $n = 186$); (d) Evergreen Needleleaf Forest plots (ENF, $n = 563$); (e) mixed forest plots ($n = 12$). PFT = Plant functional type. Canopy $N_{[\text{area}]}$ variable was log transformed to fulfil linear models assumptions.

600



Table 1. Descriptive analysis of canopy nitrogen (N) concentration ($N_{[\%]}$, $\text{g } 100\text{g}^{-1}$), foliar biomass (g m^{-2}) and canopy N content ($N_{[\text{areal}]}$, g m^{-2}) by tree species. PFT = Plant Functional Type, DBF = Deciduous Broadleaf Forest, EBF = Evergreen Broadleaf Forest, ENF = Evergreen Needleleaf Forest, mixed = mixed forest, min = minimum value, max = maximum value, mean = averaged value, sd = standard deviation, ^a codominant plots refer to the plots where two tree species were dominant in the canopy, ^b foliar biomass data was lacking for five of the plots. Foliar biomass and canopy N content statistics are thus measured on a restricted number of plots.

| Species | PFT | Number of plots | Abundance (% of total number of plots) | Canopy $N_{[\%]}$ ($\text{g } 100\text{g}^{-1}$) | | | | Foliar biomass (g m^{-2}) | | | | Canopy $N_{[\text{areal}]}$ (g m^{-2}) | | | |
|----------------------------|-------|----------------------|----------------------------------------|------------------------------------------------------|------|------|------|--------------------------------------|---------|--------|--------|---------------------------------------------------|-------|------|------|
| | | | | min | max | mean | sd | min | max | mean | sd | min | max | mean | sd |
| <i>Castanea sativa</i> | DBF | 14 | 1.7 | 1.62 | 2.81 | 2.08 | 0.36 | 18.13 | 425.90 | 203.46 | 123.49 | 0.40 | 11.99 | 4.25 | 2.89 |
| <i>Fagus sylvatica</i> | DBF | 15 | 1.8 | 1.22 | 3.13 | 2.28 | 0.61 | 49.94 | 279.86 | 173.54 | 68.70 | 1.21 | 7.40 | 3.96 | 1.95 |
| <i>Pinus halepensis</i> | ENF | 240 | 28.4 | 0.56 | 1.57 | 0.90 | 0.19 | 9.58 | 827.80 | 197.23 | 145.54 | 0.09 | 7.29 | 1.77 | 1.33 |
| <i>Pinus nigra</i> | ENF | 37 | 4.4 | 0.56 | 1.28 | 0.89 | 0.19 | 32.25 | 923.98 | 294.29 | 224.32 | 0.23 | 8.87 | 2.67 | 2.18 |
| <i>Pinus pinaster</i> | ENF | 5 | 0.6 | 0.82 | 1.08 | 0.93 | 0.13 | 271.75 | 718.87 | 501.67 | 211.53 | 2.30 | 7.69 | 4.75 | 2.25 |
| <i>Pinus pinea</i> | ENF | 5 | 0.6 | 0.75 | 1.06 | 0.95 | 0.14 | 103.28 | 275.50 | 179.74 | 66.80 | 1.08 | 2.91 | 1.71 | 0.75 |
| <i>Pinus sylvestris</i> | ENF | 198 | 23.4 | 0.67 | 2.14 | 1.11 | 0.20 | 10.48 | 828.63 | 326.44 | 181.20 | 0.10 | 12.86 | 3.65 | 2.22 |
| <i>Pinus uncinata</i> | ENF | 69 | 8.2 | 0.46 | 1.33 | 0.87 | 0.19 | 183.59 | 1744.50 | 687.22 | 345.21 | 1.41 | 16.97 | 5.92 | 3.25 |
| <i>Quercus canariensis</i> | DBF | 3 | 0.4 | 1.97 | 2.78 | 2.25 | 0.46 | 122.11 | 197.85 | 160.32 | 37.87 | 2.41 | 5.51 | 3.71 | 1.61 |
| <i>Quercus faginea</i> | DBF | 4 | 0.5 | 1.49 | 2.11 | 1.82 | 0.31 | 10.34 | 419.14 | 233.47 | 187.01 | 0.17 | 8.83 | 4.64 | 4.09 |
| <i>Quercus humilis</i> | DBF | 9 | 1.1 | 1.53 | 3.11 | 2.41 | 0.42 | 56.12 | 337.33 | 142.65 | 92.11 | 1.21 | 8.64 | 3.33 | 2.19 |
| <i>Quercus cerrides</i> | DBF | 17 | 2.0 | 1.44 | 2.80 | 2.07 | 0.37 | 12.97 | 834.68 | 262.24 | 237.49 | 0.29 | 15.42 | 5.06 | 4.31 |
| <i>Quercus ilex</i> | EBF | 160 | 18.9 | 0.81 | 2.87 | 1.32 | 0.26 | 16.63 | 1033.31 | 378.23 | 238.61 | 0.22 | 16.61 | 4.95 | 3.23 |
| <i>Quercus petraea</i> | DBF | 17 | 2.0 | 1.37 | 2.70 | 2.21 | 0.41 | 20.45 | 741.42 | 279.96 | 229.78 | 0.32 | 15.37 | 5.98 | 4.66 |
| <i>Quercus suber</i> | EBF | 23 | 2.7 | 1.25 | 2.08 | 1.55 | 0.21 | 26.26 | 219.05 | 110.49 | 55.65 | 0.40 | 4.34 | 1.72 | 0.96 |
| Codominant ^a | mixed | 30 (25) ^b | 3.5 | 0.92 | 2.54 | 1.45 | 0.41 | 23.45 | 342.58 | 153.70 | 77.39 | 0.33 | 5.74 | 2.06 | 1.02 |



Table 2. Observed linear regression equations between the MERIS Terrestrial Chlorophyll Index (MTCI) (-) and canopy nitrogen concentration (CN_[%], %N) for different subgroups. Number of plots (n), determination coefficient (r²) and p-value. PFT = Plant Functional type, DBF = Deciduous Broadleaf Forest, EBF = Evergreen Broadleaf Forest, ENF = Evergreen Needleleaf Forest, mixed = mixed forest.

| group | n | linear regression | 95% confidence interval intercept | 95% confidence interval slope | r ² | p-value |
|---------|-----|-------------------------------|--------------------------------------|----------------------------------|----------------|---------|
| overall | 846 | MTCI = 2.18 + 0.79 log(CN[%]) | [2.148, 2.202] | [0.713, 0.867] | 0.32 | < 0.000 |
| DBF | 80 | MTCI = 2.07 + 0.95 log(CN[%]) | [1.782, 2.359] | [0.585, 1.321] | 0.25 | < 0.000 |
| EBF | 186 | MTCI = 2.39 + 0.29 log(CN[%]) | [2.311, 2.477] | [0.044, 0.535] | 0.03 | 0.021 |
| ENF | 564 | MTCI = 2.13 + 0.61 log(CN[%]) | [2.100, 2.169] | [0.463, 0.762] | 0.10 | < 0.000 |
| mixed | 16 | MTCI = 2.05 + 1.35 log(CN[%]) | [1.634, 2.463] | [0.534, 2.171] | 0.47 | 0.003 |



Table 3. Observed linear regressions equations between the MERIS Terrestrial Chlorophyll Index (MTCI) (-) and canopy nitrogen content ($CN_{[area]}$, $g\ m^{-2}$) for different subgroups. Number of plots (n), determination coefficient (r^2) and p-value. PFT = Plant Functional type, DBF = Deciduous Broadleaf Forest, EBF = Evergreen Broadleaf Forest, ENF = Evergreen Needleleaf Forest, mixed = mixed forest.

| group | n | linear regression | 95% confidence interval intercept | 95% confidence interval slope | r^2 | p-value |
|---------|-----|----------------------------------------|-----------------------------------|-------------------------------|-------|---------|
| Overall | 841 | $MTCI = 2.08 + 0.20 \log(CN_{[area]})$ | [2.036, 2.116] | [0.172, 0.234] | 0.17 | <0.000 |
| DBF | 80 | $MTCI = 2.72 + 0.06 \log(CN_{[area]})$ | [2.576, 2.872] | [-0.043, 0.154] | 0.02 | 0.263 |
| EBF | 186 | $MTCI = 2.39 + 0.07 \log(CN_{[area]})$ | [2.315, 2.464] | [0.023, 0.123] | 0.04 | 0.005 |
| ENF | 563 | $MTCI = 1.94 + 0.20 \log(CN_{[area]})$ | [1.907, 1.989] | [0.168, 0.236] | 0.20 | <0.000 |
| mixed | 12 | $MTCI = 2.43 + 0.34 \log(CN_{[area]})$ | [2.045, 2.815] | [-0.263, 0.949] | 0.14 | 0.236 |



11 Appendix




Figure A 1. Allometric relationship between the branch diameter (mm) and the foliar dry mass (g) for *Pinus sylvestris* (n = 208). This is an example of the relationships used to obtain the foliar biomass data of the forest plots. Modified from (Gracia et al., 2004).




Figure A 2. The upper right part of this figure shows the Pearson correlation matrix between canopy $N_{[%N]}$ (%N), canopy $N_{[area]}$ ($g\ m^{-2}$) and foliar biomass ($g\ m^{-2}$) variables for deciduous broadleaf forest plots (DBF), $n = 80$. The diagonal presents the histogram of the variable on the x-axis, while the y-axis represents the number of counts. The lower left part of this figure represents the scatterplots between the variables.




Figure A 3. The upper right part of this figure shows the Pearson correlation matrix between canopy N_[%N] (%N), canopy N_[area] (g m⁻²) and foliar biomass (g m⁻²) variables for evergreen broadleaf forest (EBF) plots, n = 186. The diagonal presents the histogram of the variable on the x-axis, while the y-axis represents the number of counts. The lower left part of this figure represents the scatterplots between the variables.




Figure A 4. The upper right part of this figure shows the Pearson correlation matrix between canopy N [%N], canopy N [area] (g m⁻²) and foliar biomass (g m⁻²) variables for evergreen needleleaf forest (ENF) plots, $n = 563$. The diagonal presents the histogram of the variable on the x-axis, while the y-axis represents the number of counts. The lower left part of this figure represents the scatterplots between the variables.