

COMMENTARY

Open Access



A better use of fertilizers is needed for global food security and environmental sustainability

Josep Penuelas^{1,2*} , Fernando Coello^{1,2} and Jordi Sardans^{1,2}

Abstract

The massive use of fertilizers during the last decades allowed a great increase in the global capacity of food production. However, in the last years, several studies highlight the inefficiency and country asymmetries in the use of these fertilizers that generated environmental problems, soil nutritional imbalances and not optimal food production. We have aimed to summarize this information and identify and disentangle the key caveats that should be solved. Inadequate global management of fertilization produces areas with serious nutrient deficits in croplands linked with insufficient access to fertilizers that clearly limit food production, and areas that are overfertilized with the consequent problems of environmental pollution affecting human health. A more efficient use of nitrogen (N), phosphorus (P) and potassium (K) fertilizers for food security while preserving the environment is thus needed. Nutrient imbalances, particularly the disequilibrium of the N:P ratio due to the unbalanced release of N and P from anthropogenic activities, mainly by crop fertilization and expanding N-fixing crops that have continuously increased the soil N:P ratio, is another issue to resolve. This imbalance has already affected several terrestrial and aquatic ecosystems, altering their species composition and functionality and threatening global biodiversity. The different economic and geopolitical traits of these three main macronutrient fertilizers must be considered. P has the fewest reserves, depending mostly on mineable efforts, with most of the reserves concentrated in very few countries (85% in Morocco). This problem is a great concern for the current and near-future access to P for low-income countries. N is instead readily available due to the well-established and relatively low-cost Haber–Bosch synthesis of ammonium from atmospheric N₂, which is increasingly used, even in some low-income countries producing an increasing imbalance in nutrient ratios with the application of P and K fertilizers. The anthropogenic inputs of these three macronutrients to the environment have reached the levels of the natural fluxes, thereby substantially altering their global cycles. The case of the excess of N fertilization is especially paradigmatic in several areas of the world, where continental water sources have become useless due to the higher nitrate concentrations. The management of N, P and K fertilizers is thus in the center of the main dichotomy between food security and environmentally driven problems, such as climate change or eutrophication/pollution. Such a key role demands new legislation for adopting the well-known and common-sense 4R principle (right nutrient source at the right rate, right time and right place) that would help to ensure the appropriate use of nutrient resources and the optimization of productivity.

Keywords Fertilizers, Food security, Environmental security, Nitrogen, Phosphorus, Potassium, Imbalance, Stoichiometry, Eutrophication, Global cycles

*Correspondence:

Josep Penuelas

josep.penuelas@uab.cat

¹ CSIC, Global Ecology Unit CREA-FC-SCIC-UAB, 08913 Bellaterra, Catalonia, Spain

² CREA, 08913 Cerdanyola del Vallès, Catalonia, Spain

Introduction

The global population is expected to increase by about 35% over the next 40 years [87]. Agricultural output will need to increase substantially to accommodate the growing population. Most of the increase (in agricultural



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

output) is expected to be from producing more food on existing farmland (i.e., intensification), although some new farmland will likely be needed [26]. Such intensification and expansion might, however, lead to undesirable impacts on carbon (C) stocks in soil and vegetation and on biodiversity in the most productive croplands of the world [8].

Boosting crop yields and closing the gap between actual and attainable yields can be achieved by implementing and advancing numerous practices and technologies, e.g., the adequate use of fertilizers and efficient nutrient management can play key roles for global food security [82, 89]. However, the fertilization intensification of the last decades aimed to increase yields has produced some new global environmental and geopolitical problems, such as nutrient imbalances, [31, 50, 59, 61, 68], leaching of nutrients from crops to environment and the associated impacts [45, 68, 72, 85, 94], and increasing cost of fertilizers with serious geopolitical and economic problems for the food security in poor countries [3, 40, 57, 89]. The rise of fertilizer application at global scales has grown exponentially for N than for P and K (Lu and Tian [50]), [59, 60, 61, 75]. The levels of human driven N, P and K annual loads to environment have reached similar values than natural fluxes and a substantial rise in human driven N:P and N:K ratios [59, 60, 61, 72, 75].

In this commentary, we aim to discuss the problems caused and associated by the exponential of fertilizers use such as environmental pollution, global nutrient imbalances and food security mainly in poor countries, their causes, interactions, and consequences, both for current conditions and for the coming decades, also disentangling the potential solutions to mitigate and adapt to this global change driver, the human driven global changes in N, P and K stocks, cycling and stoichiometry produced by continuous temporally and spatially asymmetric consumption of fertilizers. This should improve a global view linking human use of fertilizers, with global food security and environmental quality and environmental services potential. We aimed to make so by summarizing and connecting the main bibliography on this topic.

We have checked the current bibliography and reports about global use of fertilizers and their relationships and impacts on food production and environment. We have checked google, google scholar and Web of Science. We have used combinations of words, such as: “fertilizer & environment”, “fertilizer & food security”, “fertilizer & phosphorus”, “fertilizer & nitrogen”, “fertilizer & potassium” “N:P ratio & fertilizer & environment”, “N:P ratio & phosphorus & nitrogen & fertilizer & environment”, “environment & food security & poor countries”, “nitrogen & phosphorus & N: ratio”, “fertilizer & phosphorus & nitrogen”, “fertilizer & phosphorus & nitrogen &

potassium”, “ We have thereafter selected the studies of the last 30 years with a global perspective.

N fertilization

N fertilization has increased by an order of magnitude in the last six decades [66]. Converting atmospheric N₂ to ammonium as a source for N fertilization has been the main process responsible for the increased agricultural yield since the end of World War II [24]. The average increase in yield during 1930–2000 attributable to inputs of N fertilizers generally ranged from about 40–64% in temperate climates (USA and England) and tended to be much higher in the tropics [24, 82]. Higher N inputs to improve crop yield should be possible without exceeding the critical regional N concentration in runoff in some areas of the world, such as southeastern Asia, Latin America, Oceania, the Caribbean and sub-Saharan Africa (except South Africa) [8].

The response of agricultural systems to increased N fertilization has evolved differently in countries during the last five decades. Some countries have improved their agro-environmental performances, but increased fertilization in other countries has produced low agronomic benefits and higher environmental losses [43], such as accelerated emissions of nitrous oxide leading to global climate change and high N loadings leading to contaminated drinking water and toxic algal blooms in downstream ecosystems [17, 30]. In addition to the strictly environmental impact, excess N fertilization has also been associated with threats to human health, such as cancer, upper respiratory diseases [101] and immunogenic pathologies, such as coeliac disease [59].

Achieving a worldwide reduction of total anthropogenic inputs of N into the biosphere and humans is thus a major challenge that requires sustained actions to improve the management practices of N fertilization and reduce the losses of N to the environment. More N, though, will be needed to feed the additional 1.5–2 billion people that will be added to the global population before population growth stabilizes later in this century [46]. Some studies have observed that an adequate management of N fertilizers in several countries has influenced N pollution much more than crop yields and that the trade-off between reducing N pollution and increasing yields was small. Countries that have caused 35% less N pollution than their neighbors generally only had a 1% loss of potential yield. Explanatory variables of which countries cause the most pollution relative to their crop yields include economic development, population size, institutional quality and foreign financial flows to land resources and overall agricultural intensity and share in the economy. These findings provide consistent evidence that many national governments have an impressive

capacity to reduce global N pollution without having to sacrifice much agricultural production [97]. Unique measures could likely resolve this complex issue, and a combination of measures should probably be immediately applied at the global scale.

Several options for reducing the environmental effects of the food system while improving world capacity to supply food to humans have been proposed. Their synergistic combination will be needed to sufficiently mitigate the projected increase in environmental pressures [8]. These options include dietary changes toward healthier, more plant-based diets, improvements in technologies and management and reductions in the loss and waste of food [12, 37]. However, scenario projections indicate that supply-side measures such as the improvement of N-use efficiency are more important than demand-side efforts for food security when introducing regional N targets. International trade also plays a key role in sustaining global food security under N-boundary constraints if only a limited set of mitigative options is deployed [8].

Agricultural sustainability, defined as the ability to use soil crops to produce continuous food without environmental (basically soil) degradation and other environmental impacts [99], proposes a large list of new or changed practices that can be applied including the inoculation of Arbuscular mycorrhizal fungi (AMF), cyanobacteria, and beneficial nematodes, which enhance water use efficiency and nutrient availability to plants, phytohormones production, soil nutrient cycling, and plant resistance to environmental stresses [99]. Farming practices have shown that organic and/or regenerative farming with conservation tillage, manual and biological weeding and pest control and use of farming and crop waster as fertilizers to reduce the need to apply industrial fertilizer, smart intercrop and crop rotation managements; all them improving soil health by increasing the abundance, diversity, and activity of microorganisms [21, 33, 42, 76, 84, 91].

Precision farming has allowed to make a step forward in the technology application to improve fertilizers and other resources use efficiency and diminish environmental impacts is [29]. Precision farming aims to use the most advanced technology tools to tailor management to site crop type and environmental conditions. It is based on the “diagnostic” of the situation of crop and in base to this provide the application of adequate tools. For instance, the spectral indexes are one of the most precise diagnostic tools. The sensors and scanner are frequently mounted on tractors and provide information of soil nutrient status along crop fields, allowing a site-specific fertilization [20, 55]. Similarly, advances can be also applied in the case of N to increase its use efficiency such as those based on smart agriculture including new

fertility management that included N fertilizers with slow N releasing, sensors in soil crops to monitor the actual status of soil N, and an adequate combination of crop species in intercropping or crop rotation improving the use of N₂-fixing crops [35].

P fertilization

P fertilization has increased much less than N-fertilization in the last decades [68]. The demand for P fertilizers, however, is expected to increase in the next decades; the global peak in P production has been predicted to occur around 2030 [14]. Estimates of when existing P reserves could be exhausted, however, range widely from the next 40–400 years [15, 57], US [88]. The exact timing of peak P production is disputed, but the quality of the remaining phosphate rock is widely acknowledged to be continuously decreasing [18, 36], so production costs are increasing and thus hindering the access of low-income countries to P fertilizers. The situation can be more problematic, because some models project that the global needs of P fertilizers can be doubled by 2050 [56] and also project an increase in the loss of soil P, including croplands, by soil erosion linked to climate change, which will increase the need for P fertilizers to sustain crop production [1].

The situation is further complicated by the distribution of mineable sources of P, which are concentrated in very few countries (85% in Morocco) [57]. This unequal distribution and the unequal economic capacity of different countries has caused large surpluses in most of eastern Asia, western and southern Europe, the coastal USA and southern Brazil but deficits in Africa [52, 89]. Over-fertilized soils are frequently saturated with P because of their historically high applications [15, 60]. P deficits occur across 30% of the world’s cropland area, and prolonged P deficits can deplete soil P and limit crop yields [52]. The low-income and food-deficient countries in sub-Saharan Africa, central Asia and Latin America generally suffer from low P inputs (0–5 kg ha⁻¹) to their systems of agricultural production. P scarcity is thus seriously threatening soil fertility, agricultural production and global food security in several areas [15, 100].

Managing this scarcity, however, is possible. Up to 80% of the initial P supplied to overfertilized crops is estimated to be lost before consumption, mostly due to the erosion of soil [1, 15, 57]. Reductions in waste could free up this resource for low-income, food-deficient countries [57]. If the current volumes of P fertilizers were to be redistributed and used more efficiently at the global scale, cropland would not be deficient in P. If 21% of the P fertilizers used in all areas with high surpluses were to be redistributed across all P-deficit cropland, the total crop requirements for P in these locations would effectively be

met, eliminating all P deficits globally [52]. Opportunities for recovering P and reducing demand are thus also possible and constitute an institutional challenge [14]. To achieve P sustainability, several actors must contribute, farmers need to use P more efficiently and societies and states should develop technologies and practices to recycle P from the food chain [23]. For instance, the global increase in livestock production by threefold for human consumption over the last five decades has been a key driver of scarcity, environmental distribution, and decrease in the efficiency of P use [50]. Currently monogastric livestock, as pigs and poultry, suppose a 70% of the global livestock production, and taking into account the low capacity of these species to absorb P from phytases drive to the production of manure very rich in P and with low N:P ratios leading to a very low P use-efficiency [58, 62, 92], that, moreover, has contributed to exacerbate the environmental imbalances in N:P ratios [52, 59, 60, 77]. Thus, advance in a global change of human diet behavior toward a diet more based on plant-based foods should be considered for several reasons but also for a higher global P-use efficiency in food production [65, 96].

Furthermore, it is already possible to improve P use efficiency based on available new technologies, including the use of modern chemically modified phosphate fertilizers with controlled release of phosphorus [34] and new crop genotypes genetically engineered to be more efficient in P mobilization, capture and use [32]. All these methodological and technological advances drive to a more efficient management of N and P fertilization avoiding imbalances in N:P ratios and excessive use of fertilizers and subsequent associated cascades of food security and environmental problems [13, 63, 69, 95].

K fertilization

Fertilization with industrial fertilizers containing the third key nutrient, K, from mineable reserves has continuously increased since the Industrial Revolution. As with P, we are currently in a scenario, where rich countries tend to overfertilize with K, implying environmental problems and even potential threats to human health, whereas poor countries frequently have problems with access to K fertilizers, limiting their crop production. This dichotomy parallels a scenario of increasing limitation of mineable K sources and an increase in aridity under climate change and the potential demand by crop intensification [72]. Future access to K is thus urgently needed in regions most threatened by increasing aridity and food security (e.g., the Sahel, areas with Mediterranean climates and parts of Asia and South America) due to the crucial role of K in the uptake, transport and use efficiency of water by plants ([72], 2021). Insufficient K

supply is becoming an important constraint to crop production in developing countries [4]. This constraint can be increased by climate change, with terrestrial areas consequently becoming increasingly arid and semi-arid (Huang et al., 2016), [78]. Very low rates of application of potash fertilizer in agricultural production in India and other developing countries lead to the rapid depletion of K in the soil. The depletion of plant-available K in soils has a variety of negative impacts, including preventing optimal uses of N and P fertilizers, threatening the yields of the cropping systems and decreasing farmer income [4, 54].

The imbalances among fertilizers

This complex situation between N, P and K fertilizers leads to imbalances in their use around the world [50, 59, 60]. The case of the African continent is paradigmatic: P fertilization should be increased 2.3-fold to be optimal given the current inputs of N. Inputs of N that would allow Africa to close the gaps in yield should be increased at least fivefold, so the application of P fertilizer should be increased 11.7-fold [89]. Arranging all these imbalances in P provision would greatly increase the pressure on the current global extraction of P resources, which is not an easy solution due to the scarcity and high cost of mineable P, and/or as commented previously, with a net “transfer” of P fertilizers from overfertilized to under fertilized areas. In addition, if P fertilizers cannot be made increasingly accessible, the projections of crop yields of the Millennium Ecosystem Assessment imply an increase in nutrient deficits in developing regions [61]. In fact, some studies have found that cropland areas receiving high doses of P fertilizers have low P-use efficiencies in cropland yield production [52], and similar observations for N fertilization have been provided [2, 64]. These imbalances are a particular concern, because anthropogenic inputs of these three macronutrients to the environment have reached the levels of the natural fluxes, thereby substantially affecting their global cycles. The impacts could thus seriously affect cropland N:P:K ratios but also those of other ecosystems, both terrestrial and aquatic [59, 61, 73, 74]. The impacts of the changes in these ratios have been widely observed to alter the structure and function of several ecosystems around the world [59, 61, 73, 74].

Managing N fertilizers better could also help to resolve these general imbalances and the asymmetrical distribution of fertilizer. The maximum anthropogenic use of N fertilizers needed to prevent the substantial eutrophication of aquatic ecosystems is considered to be around 62 Tg N y^{-1} from industrial fertilizers and N-fixing crops [19], but we have already surpassed this threshold by at least threefold [81]. A few agricultural regions with very high rates of N application are the main contributors to

the transgression of this “de Vries” boundary, suggesting that a redistribution of N could simultaneously boost global crop production and reduce the transgression of the “de Vries” boundary at the regional level [81]. In the absence of N mitigation or redistribution, pressure on the environment will probably increase [79], even though the need of N fertilizers for sustaining the global human population is certainly large, even with the efficient use of the fertilizers [19]. The total amounts of N fertilizers needed to feed the human population in 2030 under the current distribution of fertilizers could reach 271 Tg N y⁻¹ [49], 4.5-fold higher than the threshold estimated by de Vries et al. [19]. Maintaining the equilibrium of the fertilizer N:P:K ratios in this scenario would require annual increases in the application P and K fertilizers that could strain the fertilizer markets even more [57], with unknown impacts on global food security, especially in low-income countries. The environmental effects of the food system in the scenario for 2050 could increase by 50–90% in the absence of technological changes and dedicated mitigative measures due to expected changes in population and income levels, reaching levels much higher than acceptable planetary boundaries that define a safe operating space for humanity [8].

The use of N, P and K fertilizers is greatly unbalanced around the world, including in the largest countries, such as India [7] and the USA [90], in all cases leading to deficient production of food [50, 68, 90]. Nutrient balance is essential for achieving global food security [6] and for conserving N and P fertilizers [53]. Lu and Tian [50] found a global increase in the fertilizer N:P ratio of 0.8 (g g⁻¹) per decade ($p < 0.05$) during 1961–2013, which may have an important global implication for anthropogenic impacts on agroecosystem functions in the long term. These imbalances can alter soil stoichiometry and have been observed to differ in important cropland areas [90]. The inexorable change in the stoichiometry of C and N relative to P has no equivalent in Earth history [61, 2020]. The N:P ratio of atmospheric deposition in several parts of the world has reached values over 100–200 on a mass basis [9]. This anthropogenic imprint on the N and P cycles and N:P stoichiometry has already had consequences in several natural ecosystems to the structure, functioning, and diversity of terrestrial [38, 59, 60, 61] and aquatic [22, 59, 60, 61, 73] organisms.

The role of the N:P ratio has been widely studied, and its importance for the growth of crop plants and for the structure, diversity and functioning of other ecosystems has also been widely observed [22, 61, 73], 2020), but the importance of N:K and P:K ratios has been studied less. Some studies, however, have observed that optimal N:K and P:K ratios in fertilizers are important for plant growth and yield in diverse crops [25, 41, 51]. N:K

and PK ratios have been associated with the trade-off between plant growth and energy production [51], and different abilities in the efficiency of energy production among N, P and K for plant growth and development have been inferred [51]. The most probable trade-off that may, at least in part, determine the optimal N:K ratio is that between the allocation to growth (where N is more determinant) versus the allocation to anti-stress defense, mainly against water stress, where the role of K is more important [44, 71, 72]. *Erica multiflora*, a sclerophyllous Mediterranean shrub, has the highest foliar N:K and P:K ratios in spring, coinciding with the period of growth, and the lowest foliar ratios in summer, coinciding with the period of water stress [67]. K fertilization, however, can also be important for plant and yield production, but only when the N supply reaches a specific level [5]. N:K ratios are larger in non-crop plant–soil systems due to N deposition [28], affecting the resistance of plants to stresses, such as drought [70]. The role of the P:K ratio in non-crop ecosystems, however, has rarely been studied. Optimal N:K and N:P ratios can be different and higher for plants under elevated atmospheric CO₂ concentrations, because higher net photosynthetic rates under elevated CO₂ concentrations increase the plant demand of all three macronutrients but with a higher proportion of N, mainly to reach a higher maximum rate of carboxylation [16]. The impacts of the unbalanced use of N and P fertilizers with K fertilizer on non-crop ecosystems, however, remains to be widely studied.

Toward a more rational use of fertilizers

The present and future management of N, P and K fertilizers at the global scale is directly linked to food security and environmental health and, therefore, to human life, health and well-being. Human food security and environmental health cannot be accomplished without measures leading to strategic and adequate N, P and K fertilization, which should be addressed simultaneously for the three nutrients.

Several strategies have been proposed to balance the N:P:K input ratios in fertilization management to optimize yield increases and the quantity and quality of production while minimizing nutrient losses and thus environmental impacts. Precision farming is thus a key tool at this regard [29]. Adoption of the 4R principle (right nutrient source at the right rate, right time and right place) should also help to ensure the appropriate use of nutrient resources and to optimize productivity [82].

New legislation considering the different fertilizers altogether is also urgently needed for achieving these objectives, and in fact, it is already in the agendas of some

high-income countries, such as the European Union [93]. We, however, need more international agreement about a global strategy of fertilization, because adequate fertilizer use increases water-use efficiency, crop and food production, the resistance of crop production against aridity and the preservation of environments and biodiversity and can provide several feedbacks to mitigate climate change [59, 60, 98]. In this context and at this moment, the European Union is a paradigm; in fact, European Sustainable Phosphorus Platform for an Integrated Nutrient Management Action Plan (INMAP) is fully in line with these ambitions and its pressure on European Commission in principle will act to reduce nutrient losses by at least 50%, while ensuring that there is no deterioration in soil fertility, with a short-time scenario of a reduction of the use of fertilizers by at least 20% by 2030 [93]. Unfortunately, apart from general declarations, concrete law measures to more controlled, efficient and environmentally friendly use of fertilizers are at this moment effective in very few other parts of world. Recently, in Canada the government has proposed a reduction of a 30% in the use of industrial fertilizers within the global country plan to reduce greenhouse emissions [27]. Theoretically, China is another state with strict legislation of fertilizers use [11], being prohibited to import, produce, sell or use un-registered fertilizers, and fertilizer sold in China shall also meet relevant product standards. However, there is still a lot of room in most countries to take a more serious policy about fertilizer use and to endorse all the technical improvements to reduce fertilizer use while increasing use efficiency [10, 83, 89]. It is necessary to overcome the constraints imposed by conventional intensive agriculture and the scarce economical capacity and little access of farmers of several countries to adequate information [10, 83, 89].

Conclusions

Human-driven N and P loads to environment have reached levels comparable to natural fluxes. The continuous exponential increases of N loads rising faster than P and K loads have increased N:P ratios in several areas of the world with unprecedented impacts on ecosystems and their services capacities. These imbalances have serious impacts on food production capacity, mainly in poor countries, where the increases in N:P and N:K ratios are further favoured by the higher costs of K and mainly P fertilizers than the costs of N fertilizers. In the case of the excess of N, its consequences can also affect human health by the pollution of waters and excessive N fertilization. Aridity is expected to increase in the next decades and more lands will be affected by decreased availability of water. Future access to K is thus urgently needed in regions most threatened by this

increasing aridity (e.g., the Sahel, areas with Mediterranean climates and parts of Asia and South America) due to the crucial role of K in the uptake, transport and use efficiency of water by plants, that makes K a necessary tool to try to maintain acceptable levels of food production under increasing water scarcity. In the case of P, if it is sufficiently applied, it can be stored in soil in not available forms, whereas mine reserves can be emptied faster and its access can progressively become prohibitive for poor countries. This can, moreover, be related to higher prices of P fertilizers which can hinder even more the access to fertilizer of millions of farmers in poor countries and decrease even more the already low efficiency of crop production in several areas of poor countries, such as in Africa or Asia impairing even more their food security. There are though several new technologies and crop management methodologies that can help to improve fertilizer use and efficiency. Smart, precision and regenerative agriculture approaches together with new biotechnologies application can help to correct this global change driver, so they should be quickly at global scale. Meanwhile, new legislation adopting the well-known and common-sense 4R principle (right nutrient source at the right rate, right time and right place) would help to ensure the appropriate use of fertilizers and the optimization of crop productivity for food security and environmental sustainability.

Acknowledgements

We thank the insightful exchange of ideas on the topic of this article with the members of the Global Ecology Unit CREA-FCM-CSIC-UAB, Barcelona, Catalonia, Spain.

Author contributions

JP and JS designed the study, JP, FC and JS gathered the information and wrote this comment. All authors read and approved the final manuscript.

Funding

Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. Authors acknowledge the financial support from the Catalan Government Grant SGR 2017–1005, the Spanish Government Grants PID2019-110521 GB-I00, PID2020115770RB-I, and TED2021-132627B-I00, and the Fundación Ramón Areces grant CIVP20A6621.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

We approve and consent.

Consent for publication

We consent.

Competing interests

Authors declare no competing interests.

Received: 29 September 2022 Accepted: 31 January 2023
Published online: 23 March 2023

References

- Alewell C, Ringeval B, Ballabio C, Robinson DA, Panagos P, Borrelli P. Global phosphorus shortage will be aggravated by soil erosion. *Nature Com.* 2020;11:4546.
- Albornoz F. Crop responses to nitrogen overfertilization: a review. *Scientia Horticult.* 2016;205:79–83.
- Bonilla-Cedrez C, Chamberlin J, Hijmans RJ. Fertilizer and grain process constrain food production in sub-Saharan Africa. *Nature Food.* 2021;2(2):766–72.
- Cakmak I. Potassium for better crop production and quality. *Plant Soil.* 2010;2010(335):1–2.
- Cardoso DSCP, Sedyama MAN, Poltronieri Y, Fonseca MCM, Neve YF. Effect of concentration and N: K ratio in nutrient solution for hydroponic production of cucumber. *Rev Caatinga.* 2017;30:818–24.
- Ciampitti IA, Vyn TJ. Understanding global and historical nutrient use efficiencies for closing maize yield gaps. *Agr J.* 2014;2014(106):2107–17.
- Chand R, Pavithra S. Fertilizer use and imbalances in India. *Econ Pol Wkly.* 2015;1:98–104.
- Chan J, Havlik P, Leclère D, de Vries W, Valin H, Deppermann A, Hasegawa T, Obersteiner M. Reconciling regional nitrogen boundaries with global food security. *Nature Food.* 2021;2:700–11.
- Chen HY, Huang LM, Ho TY, Chiang KP, Chou WC. A study of the nitrogen and phosphorus imbalance in East Asia based on the distribution patterns of an stoichiometry variation in global atmospheric nitrogen and phosphorus. *Atm Env.* 2021;266: 118691.
- Chojnacka K, Moustakes K, Witek-Krowiak A. Bio-based fertilizers: a practical approach towards circular economy. *Bior Technol.* 2020;295: 122223.
- CIRS. 2022. www.cirs-group.com/en/agrochemicals/overview-of-fertilizer-regulations-in-china.
- Clark M, Hill J, Tilman D. The diet, health and environmental trilemma. *An Rev Env Resour.* 2018;43:109–34.
- Cordell D, Schmid Neset TS, Prior T. The phosphorus mass balance: identifying “hotspots” in the food system as a roadmap to phosphorus security. *Cur Opin Biotechnol.* 2012;23:839–45.
- Cordell D, Drangert JO, White S. The story of phosphorus: global food security and food for thought. *Glob Env Change.* 2009;19:292–305.
- Cordell D, White S. Peak phosphorus: clarifying the key issues of a vigorous debate about long-term phosphorus security. *Sustainability.* 2011;3:2027–49.
- Dang QL, Li J, Man R. N/P/K ratios and CO₂ concentration change nitrogen photosynthesis relationships in black spruce. *Am J Plant Sci.* 2021;12:1090–105.
- Davidson EA. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geosci.* 2009;2:659–62.
- Desmidt E, Ghyselbrecht K, Zhang Y, Pinoy L, Van der Brugge B, Verstraete W, Rabaey K, Meesschaert B. Global phosphorus scarcity and full-scale P-recovery techniques: a review. *Crit Rev Env Sci Technol.* 2014;45:336–84.
- De Vries W, Kros J, Kroeze C, Seitzinger SP. Assessing planetary and regional nitrogen boundaries related to food security and adverse environmental impacts. *Cur Opin Env Sustain.* 2013;5:392–402.
- Diacono M, Rubino P, Montemurro F. Precision nitrogen management of wheat. A review *Agron Sustain Dev.* 2013;33:219–41.
- Dong K, Dong Y, Zheng L, Tang Z, Yang X. Faba bean fusarium wilt (*Fusarium oxysporum*) control and its mechanism in different wheat varieties and faba bean intercropping system. *Chin J Appl Ecol.* 2014;25:1979–87.
- Elser JJ, Andersen T, Baron JS, Bergström AK, Jansson M, Kyle M, Nydick KR, Steger Hessen DO. Shifts in lake N: P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition. *Science.* 2009;326:835–7.
- Elser JJ. Phosphorus: a limiting nutrient for humanity? *Cur Opin Biotechnol.* 2012;23:833–8.
- Erisman JW, Sutton MA, Galloway J, Klimont Z, Winiwarter W. How a century of ammonia synthesis changed the world. *Nature Geosci.* 2008;1:636–9.
- Fallah M, Delshad M, Sheikhi H. The effects of cluster pruning and the N: K ratio on greenhouse tomato yield quality. *Horticult Env Biotechnol.* 2021;6:691–700.
- FAO. The future of food and agriculture. Rome: Trends and Changes; 2017.
- Farm Journal, 2022. www.agweb.com/news/policy/politics/new-policy-forces-canadian-producers-cut-back-fertilizer.
- Ferretti M, Calderisi M, Marchetto A, Waldner P, THimonier A, Jonard M, Cools N, Rautio P, Clarke N, Hansen K, Merilä P, Potocic N. Variables related to nitrogen deposition improve defoliation models for European forests. *An For Sci.* 2015;72(897):906.
- Finger R, Swinton SM, El Benni N, Walter A. Precision farming at the nexus of agricultural production and the environment an. *Rev Res Econ.* 2019;11:313–35.
- Galloway JN, Aber JD, Erisman JW, Seitzinger SP, Howarth RW, Cowling EB, Cosby BJ. The nitrogen cascade. *Bioscience.* 2003;53:341–56.
- MacDonald GK, Bennett EM, Potter PA, Ramankutty N. Agronomic phosphorus imbalances across the world's croplands. *PNAS.* 2021;108:3086–91.
- Gaxiola RA, Edwards M, Elser JJ. A transgenic approach to enhance phosphorus use efficiency in crops as part of a comprehensive strategy for sustainable agriculture. *Chemosphere.* 2011;84:840–5.
- Govaerts B, Fuentes M, Mezzalama M, Nicol JM, Deckers J, Etchevers JD, Figueroa-Sandoval B, Sayre KD. Infiltration, soil moisture, root rot and nematode populations after 12 years of different tillage, residue and crop rotation managements. *Soil Tillage Res.* 2007;94:209–19.
- Guelfi D, Nunes APP, Fernandes Sakis L, Oliveira DP. Innovative phosphate fertilizer technologies to improve use efficiency in agriculture. *Sustain.* 2022;14:14266.
- Herrera JM, Rubio G, Häner LL, Delgado JA, Lucho-Constantino Islas-Valdez S, Pellet D. Emerging and established technologies to increase nitrogen use efficiency of cereals. *Agron.* 2016;6:25.
- Herrera-Estrella L, López-Arredondo D. Phosphorus: the underrated element for feeding the world. *Trends Plant Sci.* 2016;21:461–3.
- Houlton BZ, Almaraz M, Aneja V, Austin AT, Bai E, Cassman KG, Gu B, Yao G, Matinelli LA, Scow K, Schlesinger WH, Tomich TP, Wang C, Hang X. A world of cobenefits: solving the global nitrogen challenge. *Earth's Future.* 2019;7:865–72.
- Huang WJ, Zhou GY, Liu JX. Nitrogen and phosphorus status and their influence on aboveground production under increasing nitrogen deposition in three successional forest. *Acta Oecol.* 2012;44:20–7.
- Huang J, Ji M, Xie Y, Wang S, He Y, Ran J. Global semi-arid climate change over last 60 years. *Clim Dyn.* 2017;46:1131–50.
- Khavarov N, Obersteiner M. Global phosphorus fertilizer market and national policies: a case study revisiting the 2008 price peak. *Front Nutr.* 2017;4:art22.
- Kraus HT, Warren SL, Bjorkquist GJ, Lowder AW, Tchir CM, Walton KN. Nitrogen:phosphorus:potassium ratios affect production of two herbaceous perennials. *HostScience.* 2011;2016(46):776–83.
- Larkin RP, Honeycutt CW, Olanya OM, Halloran JM, He Z. Impacts of crop rotation and irrigation on soilborne diseases and soil microbial communities. In: He Z, Larkin R, Honeycutt W, editors. *Sustainable Potato Production: Global Case Studies*. Dordrecht: Springer; 2012.
- Lassaletta L, Billen G, Grizzetti B, Anglade J, Garnier J. 50 year trends in nitrogen use efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland. *Env Res Let.* 2014;9: 105011.
- Lawniczak AE, Gusewell S, Verhoeven JTA. Effect of N: K supply ratios on the performance of three grass species from herbaceous wetlands. *Basic Appl Ecol.* 2009;2009(10):715–25.
- Lee EK, Zhang X, Adler PR, Kleppel GS, Romeiko XX. Spatial and temporally explicit life cycle global warming, eutrophication, and acidification impacts from crop production in the US midwest. *J Clean Produc.* 2020;242:118645.
- Leite JC, Caldeira S, Watzl B, Wollgast J. Healthy low nitrogen footprint diets. *Glob Food Sec.* 2020;24: 100342.

47. Lissbrant S, Brouder SM, Cunningham SM, Volenec JJ. Identification of fertility regimes that enhance long-term productivity of alfalfa using cluster analysis. *Agronomy J*. 2020;102:580–91.
48. Liu Y, Pan X, Li AJ. 1961–2010 record of fertilizer use, pesticide application and cereal yields: a review. *Agron Sustain Dev*. 2015;35:83–93.
49. Liu J, Ma K, Ciais P, Polasky S. Reducing human nitrogen use for food production. *Sci Rep*. 2016;6:30104.
50. Lu C, Tian H. Global nitrogen and phosphorus fertilizer use for agriculture production in the past half century: shifted hot spots and nutrient imbalance. *Earth Syst Sci Data*. 2017;9:181–92.
51. Ma J, Chen T, Lin T, Fu E, Feng B, Li G, Li H, Li J, Wu Z, Tao L, Fu G. Functions of nitrogen, phosphorus and potassium in energy status and their influences on rice growth and development. *Rice Sci*. 2022;29:166–78.
52. MacDonald GK, Bennett EM, Potter PA, Ramankutty N. Agronomic phosphorus imbalances across the world's croplands. *Proc Natl Acad Sci*. 2011;108:3086–91.
53. Metson GS, Bennett EM, Elser JJ. The role of diet in phosphorus demand. *Env Res Lett*. 2012;7: 044043.
54. Mkangawa CZ, Mbogoni JDJ, Ley GJ, Msolla AM. 2015. Potassium for Sustainable Crop Production and Food Security: In Mkangawa CZ, Mbogoni JDJ, Ley GJ, Msolla AM (eds). *The First National Potash*. Tanzania
55. Nawar S, Corstanje R, Halcro G, Mulla D, Mouazen AM. Delineation of soil management zones for variable-rate fertilization: a review. *Adv Agron*. 2017;143:175–245.
56. Nedelciu CE, Ragnarsdottir KV, Schlyter P, Stjernquist I. Global phosphorus supply chain dynamics: assessing regional impact to 2050. *Glob Food Sec*. 2020;26: 100426.
57. Obersteiner M, Peñuelas J, Ciais P, van der Velde M, Janssens IA. The phosphorus trilemma. *Nature Geosci*. 2013;6:897–8.
58. Oster M, Reyer H, Ball E, Fornara D, McKillen J, Sorensen CU, Wimmers K. Bridging gaps in the agricultural phosphorus cycle from an animal husbandry perspective—the case of pigs and poultry. *Sustain*. 2018;10:1825.
59. Peñuelas J, Janssens IA, Ciais P, Obersteiner M, Sardans J. Anthropogenic global shifts in biospheric N and P concentrations and ratios and their impacts on biodiversity, ecosystem productivity, food security, and human health. *Glob Change Biol*. 2020;26(1962):1985.
60. Peñuelas J, Gargallo-Garriga A, Janssens IA, Ciais P, Obersteiner M, Klem K, Urban O, Zhu YG, Sardans J. Could global intensification of nitrogen fertilization increase immunogenic proteins and favour the spread of coeliac pathology? *Foods*. 2020;9:1602.
61. Peñuelas J, Poulter B, Sardans J, Ciais P, van der Velde M, Bopp L, Boucher O, Godderis Y, Hinsinger P, Llusia J, Nardin E, Vicca S, Obersteiner M, Janssens IA. Human induced nitrogen–phosphorus imbalances alter natural and managed ecosystems across the globe. *Nature Com*. 2013;4:2934.
62. Prasad CS, Mandal AB, Gowda NKS, Sharma K, Pattanaik AK, Tyagi PK, Elangovan AV. Enhancing phosphorus utilization for better animal production and environment sustainability. *Cur Sci*. 2015;108:1315–9.
63. Rahman S, Chowdhury RB, D'Costa NG, Milne N, Bhuiyan M, Sujauddin M. Determining the potential role of the waste sector in decoupling of phosphorus: a comprehensive review of national scale substance flow analyses. *Res Conser Rec*. 2019;144:144–57.
64. Rahman KM, Zhang D. Effects of fertilizer broadcasting on the excessive use of inorganic fertilizers and environmental sustainability. *Sustain*. 2018;10:759.
65. Reijnders L. Phosphorus resources, their depletion and conservation, a review. *Resour Conserv Recycl*. 2014;93:32–49. <https://doi.org/10.1016/j.resconrec.2014.09.006>.
100. Ringeval R, Nowak B, Nesme T, Delmas M, Pellerin S. Contribution of anthropogenic phosphorus to agricultural soil fertility and food production. *Global Biogeochem Cycles*. 2014;28:743–56.
66. Ritchie H, Roser M. Fertilizers. Published online at OurWorldInData.org. 2013. <https://ourworldindata.org/fertilizers>.
67. Rivas-Ubach A, Sardans J, Perez-Trujillo M, Estiarte M, Peñuelas J. Strong relationship between elemental stoichiometry and metabolome in plants. *Proc Natl Acad Sci USA*. 2012;109:4181–6.
68. Romero E, Ludwig W, Sadaoui M, Lassaletta L, Bouwman AF, Beusen AHW, van Apeldoorn D, Sardans J, Janssens IA, Ciais P, Obersteiner M, Peñuelas J. The mediterranean region as a paradigm of the global decoupling of N and P between soils and freshwaters. *Glob Biogeochem Cycl*. 2021;35:2020GB006874.
69. Rosemarin A, Ekane N. The governance gap surrounding phosphorus. *Nutr Cycl Agroecosyst*. 2016;104:265–79.
70. Rosengren-Brinck U, Nihlgård B. Effects of nutritional status on the drought resistance in Norway spruce. *Water Air Soil Pollut*. 1995;85:1739–44.
71. Salazar-Tortosa D, Castro J, Villar-Salvador P, Vinegla B, Matias L, Michelsen A, de Casas RR, Querejeta JI. The “Isohydric trap” a proposed feedback between water shortage, stomatal regulation, and nutrient acquisition drives differential growth and survival of European pines under climatic dryness. *Glob Change Biol*. 2018;24:4069–83.
72. Sardans J, Peñuelas J. Potassium: a neglected nutrient in global change. *Glob Ecol Biogeogr*. 2015;24:261–75.
73. Sardans J, Rivas-Ubach A, Peñuelas J. The C:N: P stoichiometry of organisms and ecosystems in a changing world: a review and perspectives. *Persp Plant Ecol*. 2012;14:33–47.
74. Sardan J, Janssens IA, Ciais P, Obersteiner M, Peñuelas J. Recent advances and future research in ecological stoichiometry. *Persp Plant Ecol Evol Syst*. 2021;50: 125611.
75. Sardans J, Peñuelas J. Potassium control of plant functions: ecological and agricultural Implications. *Plants*. 2021;10:419.
76. Shao Y, Xie Y, Wang C, Yue J, Yao Y, Liu W. Effects of different soil conservation tillage approaches on soil nutrients, water use and wheat-maize yield in rainfed dry-land regions of North China. *Eur J Agron*. 2016;81:37–45.
77. Sileshi GH, Nhamo H, Mafongoya PL, Tanimu J. (2017) Stoichiometry of animal manure and implications for nutrient cycling and agriculture in sub-Saharan Africa. *Nutr Cycl Agroecosyst*. 2017;107:91–105.
78. Spinoni J, Barbosa P, Cherlet M, Forzieri G, McCormick N, Naumann G, Vogt JV, Dosio A. How will the progressive global increase of arid areas affect population and land-use in the 21st century? *Glob Planet Change*. 2021;205:103597.
79. Springmann M, Clark M, Mason-D'Croz D, Wiebe K, Bodirsky BL, Las-saletta L, de Vries W, Vermeulen SJ, Herrero M, Carlson KM, Jonell M, Troell M, DeClerck F, Gordon LJ, Zurayk R, Scarborough P, Rayner M, Loken B, Fanzo J, Godfray H, Tilman D, Rockström J, Willet W. Options for keeping the food system within environmental limits. *Nature*. 2018;562:519–25.
80. Smil V. Nitrogen and food production: proteins for human diets. *Ambio*. 2002;31:126–31.
81. Steffen W, Richardson K, Rockström J, Cornell SE, Fetzer I, Bennett EM, Biggs R, Carpenter SR, de Vries W, de Wit CA, Folke C, Gerten D, Heinke J, Mace GM, Persson M, Ramanathan V, Reyers B, Sörlin S. Planetary boundaries: guiding human development on a changing planet. *Science*. 2015;347:736.
82. Stewart WM, Roberts TL. Food security and the role of fertilizer in supporting it. *Proc Engin*. 2012;46:76–82.
83. Stuart D, Schewe RL, McDermott M. Reducing nitrogen fertilizer application as a climate change mitigation strategy: understanding farmer decision-making and potential barriers to change in the US. *Land Use Pol*. 2013;36:210–8.
99. Tahat MM, Alananbeh KM, Othman YA, Leskovar DI. Soil health and sustainable agriculture. *Sustainability* 2020;12:4859.
84. Taiwo LB, Adebayo DT, Adebayo OS, Adediran JA. Compost and glomus mosseae for management of bacterial and fusarium wilts of tomato. *Int J Veg Sci*. 2007;13:49–61.
86. Takahashi K, Muraoka R, Otsuka K. Technology adoption, impact and extension in developing countries' agriculture: a review of recent literature. *Agric Econ*. 2019;51:31–45.
85. Tedengren M. Eutrophication and disrupted nitrogen cycle. *Ambio*. 2021;50:733–8.
101. Townsend AR. Human health effects of a changing global nitrogen cycle. *Front Ecol Environ*. 2003;1:240–6.
87. UN Population prospects 2022. <https://population.un.org/wpp/Download/Standard/Population/>.
88. US. Geological Survey. Mineral commodity summaries 2016, p. 202, U.S. Geological Survey. 2016.
89. Van der Velde M, Folberth C, Balkovic J, Ciais P, Fritz S, Janssens IA, Obersteiner M, See L, Skalsky R, Xiong W, Peñuelas J. African crop yield reductions due to increasingly unbalanced nitrogen to phosphorus consumption. *Glob Change Biol*. 2014;20:1278–88.

90. Vitousek PM, Naylor R, Crews T, David MB, Drinkwater LE, Holland E, Johnes PJ, Katzenberger J, Martinelli LA, Matson PA, Nziguhaba G, Ojima D, Palm CA, Robertson P, Sanchez PA, Townsend AR, Zhan FS. Nutrient imbalances in agricultural development. *Science*. 2009;324:1519–20.
91. Vivak K, Saharawat YS, Gathala MK, Jat AS, Singh SK, Chaudhary N, Jat ML. Effect of different tillage and seeding methods on energy use efficiency and productivity of wheat in indo-gangetic plains. *Field Crops Res*. 2013;142:1–8.
92. Wang M, Ma L, Strokal M, Chu Y, Kroeze C. Exploring nutrient Management options to increase nitrogen and phosphorus use efficiencies in food production of China. *Agric Syst*. 2018;163:58–72.
93. Wassen MJ, Schrader J, Eppinga MB, Sardans J, Berendse F, Beunen R, Peñuelas J, van Dyck J. The EU needs a nutrient directive. *Nature Rev Earth Env*. 2022;3:287–8.
94. Weihrauch C, Weber CJ. Phosphorus enrichment in floodplain subsoils as a potential source of freshwater eutrophication. *Sci Total Env*. 2020;747: 141213.
95. Weikard HP. Phosphorus recycling and food security in the long run: a conceptual modelling approach. *Food Sec*. 2016;8:405–14. <https://doi.org/10.1007/s12571-016-0551-4>.
96. Withers PJA, van Dijk KC, Neset TSS, Nesme T, Onema O, Rubaek GH, Pellerin S. Stewardship to tackle global phosphorus inefficiency: the case of Europe. *Ambio*. 2015;2015(44):S193–206.
97. Wuepper D, Le Clech S, Zilberman D, Mueller N, Finger R. Countries influence the trade-off between crop yields and nitrogen pollution. *Nature Food*. 2020;1:713–9.
98. Xiong W, Tarnavsky E. Better agronomic management increases climate resilience of maize to drought in Tanzania. *Atmosphere*. 2020;11:982.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

